arXiv:1107.1906v2 [math.AG] 11 Oct 2011

Toric Stacks I: The Theory of Stacky Fans

Anton Geraschenko and Matthew Satriano

Abstract

The purpose of this paper and its sequel [GS11b] is to introduce and develop a theory of toric stacks which encompasses and extends the notions of toric stacks defined in [Laf02, BCS05, FMN09, Iwa09, Sat09, Tyo10], as well as classical toric varieties.

In this paper, we define a *toric stack* as a quotient of a toric variety by a subgroup of its torus (we also define a generically stacky version). Any toric stack arises from a combinatorial gadget called a *stacky fan*. We develop a dictionary between the combinatorics of stacky fans and the geometry of toric stacks, stressing stacky phenomena such as canonical stacks and good moduli space morphisms.

We also show that smooth toric stacks carry a moduli interpretation extending the usual moduli interpretations of $\mathbb{P}^n$ and $[\mathbb{A}^1/\mathbb{G}_m]$. Indeed, smooth toric stacks precisely solve moduli problems specified by (generalized) effective Cartier divisors with given linear relations and given intersection relations. Smooth toric stacks therefore form a natural closure to the class of moduli problems introduced for smooth toric varieties and smooth toric DM stacks in [Cox95] and [Per08], respectively.

We include a plethora of examples to illustrate the general theory. We hope that this theory of toric stacks can serve as a companion to an introduction to stacks, in much the same way that toric varieties can serve as a companion to an introduction to schemes.

Contents

1 Introduction **2**

2 Definitions **6**
2.1 The Toric Stack of a Stacky Fan 6
2.2 The Generically Stacky Case 8

3 Morphisms of Toric Stacks **9**
3.1 Toric Morphisms are Induced by Morphisms of Stacky Fans 10

4 Fantastacks: Easy-to-Draw Examples **12**
4.1 Some Non-fantastack Examples 15

The second author is partially supported by NSF grant DMS-0943832.
2010 Mathematics Subject Classification. 14D23, 14M25
Keywords and phrases. toric, fan, stack, moduli.

5 Canonical Stacks **16**

6 Toric Good Moduli Space Morphisms **18**
6.1 Good Moduli Space Morphisms 18
6.2 Proof of Theorem 6.3 19
6.3 Examples 25

7 Moduli Interpretation of Smooth Toric Stacks **28**
7.1 Examples 30
7.2 Generically Stacky Smooth Toric Stacks 31

Appendix A: Short Exact Sequences of G_βs **32**

Appendix B: Non-uniqueness of Stacky Fans **33**
B.1 The Cohomologically Affine Case 33
B.2 Isomorphisms From Morphisms of (Generically) Stacky Fans 35

1 Introduction

Recently, a number of theories of toric stacks have been introduced [Laf02, BCS05, FMN09, Iwa09, Sat09, Tyo10]. There are several reasons one may be interested in developing such a theory. First, these stacks provide a natural place to test conjectures and develop intuition about algebraic stacks, much the same way that toric varieties do for schemes. After developing an adequate theory, they are easy to work with combinatorially, just as toric varieties are. Second, in some situations toric stacks can serve as better-behaved substitutes for toric varieties. A toric variety has a canonical overlying smooth stack. It is sometimes easier to prove results on the smooth stack and "push them down" to the toric variety. Third, with the appropriate machinery, one can show that an "abstract toric stack" (e.g. the closure of a torus within some stack of interest) often arises from a combinatorial "stacky fan." The combinatorial theory of stacky fans then allows one to effectively investigate the stack in question.

We are aware of three kinds of toric stacks in the literature.

Lafforgue's Toric Stacks. In [Laf02], Lafforgue defines a toric stack to be the stack quotient of a toric variety by its torus. These stacks are very "small" in the sense that they have a dense open point. They are rarely smooth.

Smooth Toric Stacks. Borisov, Chen, and Smith defined smooth toric Deligne-Mumford stacks in [BCS05]. These are the stacks studied in [FMN09] and [Iwa09]. They are smooth and have simplicial toric varieties as their coarse moduli spaces. The second author generalized this approach in [Sat09] to include certain smooth toric Artin stacks which have toric varieties as their good moduli spaces.

Toric Varieties and Singular Toric Stacks. Toric varieties are neither "small" nor smooth in general, so the standard theory of toric varieties is not subsumed by the above approaches. In

[Tyo10, §4], Tyomkin introduces a definition of toric stacks which includes all toric varieties (in particular, they may be singular). However, these toric stacks always have finite diagonal, and the definition is "very local."

In this paper we propose a definition of toric stack that encompasses all of the above approaches. Before giving precise definitions, let us give a brief overview of the theory we propose and the contents of the two papers Toric Stacks I and II (this paper and [GS11b]). We define a *toric stack* to be any quotient $\mathcal{X} = [X/G]$ where X is a normal toric variety and G is a subgroup of the torus T of X. In analogy with toric varieties, we show:

1. Every such toric stack $\mathcal{X}$ comes from a combinatorial gadget, which we call a stacky fan. There is a rich dictionary between the combinatorics of stacky fans and the geometry of toric stacks.
2. With suitable hypotheses (see the theorem below), if $\mathcal{X}$ is any stack with a dense torus T such that the action of T on itself extends to $\mathcal{X}$, then $\mathcal{X}$ is a toric stack.

These two points are the subjects of Toric Stacks I and II, respectively. In this paper, we define a notion of stacky fan, show that every toric stack comes from a stacky fan, and develop the correspondence between stacky fans and their associated toric stacks, thereby allowing one to easily read off properties of a toric stack from its stacky fan. In Toric Stacks II, we prove

Theorem ([GS11b, Theorem 6.1])**.** *Let $\mathcal{X}$ be an Artin stack over an algebraically closed field k of characteristic 0. Suppose $\mathcal{X}$ has an action of a torus T and a dense open substack which is T-equivariantly isomorphic to T. Then $\mathcal{X}$ is a toric stack if and only if the following conditions hold:*

1. *$\mathcal{X}$ is normal, reduced, and of finite type,*
2. *$\mathcal{X}$ has affine diagonal,*
3. *geometric points of $\mathcal{X}$ have linearly reductive stabilizers, and*
4. *every point of $[\mathcal{X}/T]$ is in the image of an étale representable map from a stack of the form $[U/G]$, where U is quasi-affine and G is an affine group.*

Let us now describe in detail the contents of this paper. Above, we defined a toric stack to be the stack quotient of a normal toric variety X by a subgroup G of its torus T_0. Note that the stack $[X/G]$ has a dense open torus $T = T_0/G$ which acts on $[X/G]$. Many of the existing theories of toric stacks allow for generic stackiness; if our theory is to encompass all of the previous ones, we must also introduce a notion of "generically stacky toric stack". We do as follows:

An integral T-invariant substack of $[X/G]$ is necessarily of the form $[Z/G]$ where $Z \subseteq X$ is an integral T_0-invariant subvariety of X.[1] The subvariety Z is naturally a toric variety whose torus T' is a quotient of T_0. The quotient stack $[Z/G]$ contains a dense open "stacky torus" $[T'/G]$ which acts on $[Z/G]$.

---

[1] Note that Z must be irreducible because G cannot permute the irreducible T_0-invariant subvarieties of X.

Definition 1.1. In the notation of the above two paragraphs, a *toric stack* is an Artin stack of the form $[X/G]$, together with the action of the torus $T = T_0/G$. A *generically stacky toric stack* is an Artin stack which is isomorphic to a closed torus-invariant substack of a toric stack, i.e. is of the form $[Z/G]$, together with the action of the stacky torus $[T'/G]$.

This definition encompasses and extends the three kinds of toric stacks listed above:

- Taking G to be trivial, we see that any toric variety X is a toric stack.
- Smooth toric Deligne-Mumford stacks in the sense of [BCS05, FMN09, Iwa09] are smooth generically stacky toric stacks which happen to be separated and Deligne-Mumford. This is explained in the discussion immediately after Definition 2.19.
- Toric stacks in the sense of [Laf02] are toric stacks that have a dense open point (i.e. toric stacks for which $G = T_0$).
- A toric Artin stacks in the sense of [Sat09] is a smooth generically stacky toric stack with finite generic stabilizer and a toric variety of the same dimension as a good moduli space. See Sections 4 and 6.
- Tyomkin's toric stacks are generically stacky toric stacks as well, but this fact requires some theory. See [GS11b, Remark 6.3].

Just as toric varieties can be understood in terms of fans, toric stacks can be understood in terms of combinatorial objects called *stacky fans*, which are defined in Section 2. Once the definitions are in place, it will be clear that any morphism of stacky fans induces a toric morphism of the corresponding toric stacks. In Section 3, we prove that the converse is true as well: any toric morphism of toric stacks is induced by a morphism of stacky fans (Theorem 3.4).

Sections 2 and 3 provide a sufficient base to generate interesting examples. In Section 4, we highlight a particularly easy to handle class of toric stacks, which we call *fantastacks*. The stacks defined in [BCS05] and [Sat09] which have no generic stabilizer are fantastacks. Generically stacky toric stacks are considerably more general than fantastacks, but it is sometimes easiest to understand a generically stacky toric stack in terms of its relation to some fantastack. For example, the stacks defined in [BCS05] and [Sat09] are closed substacks of fantastacks.

$$\{\text{fantastacks}\} \subset \{\text{toric stacks}\} \subset \{\text{generically stacky toric stacks}\}$$

Section 5 is devoted to the construction of the *canonical stack* over a toric stack. The main result is Proposition 5.7, which justifies the terminology by showing that canonical stacks have a universal property. Canonical stacks are minimal "stacky resolutions" of singularities. Heuristically, the existence of such a resolution is desirable because it is sometimes possible to prove theorems on the smooth resolution and then descend them to the singular base. Indeed, the main theorem of Toric Stacks II, [GS11b, Theorem 6.1], is proved in this way.

In Section 6, we classify toric good moduli space morphisms in the sense of [Alp08]. Good moduli space morphisms generalize the notion of a coarse moduli space and that of a good quotient in the sense of [GIT]. Good moduli space morphisms are of central interest in the theory of moduli, so it is useful to have tools for easily identifying and handling many examples.

In Section 7, we prove a moduli interpretation for smooth toric stacks (Theorem 7.7). That is, we characterize morphisms to smooth toric stacks from an arbitrary source, rather than only toric morphisms from toric stacks. The familiar moduli interpretation of $\mathbb{P}^n$ is that specifying a morphism to $\mathbb{P}^n$ is equivalent to specifying a line bundle, together with $n+1$ sections that generate it. Cox generalized this interpretation to smooth toric varieties in [Cox95], and Perroni further generalized it to smooth toric Deligne-Mumford stacks in [Per08]. Smooth toric stacks are the natural closure of this class of moduli problems. In other words, any moduli problem of the same sort as described by Cox and Perroni is represented by a smooth toric stack (see Remark 7.10).

Just as toric varieties can be defined over an arbitrary base, much of this paper can be done over an arbitrary base, but we work over an algebraically closed field in order to avoid imposing confusing hypotheses (e.g. every subgroup of a torus we consider is required to be diagonalizable). A stacky fan defines a toric stack over an arbitrary base, and morphisms of stacky fans induce morphisms of toric stacks. However, if the base is disconnected, not every toric morphism is induced by a morphism of stacky fans.

Remark 1.2 (The Log Geometric Approach). There is yet another approach to toric geometry, namely that of log geometry. In this paper, we do not develop this approach to toric stacks. We refer the interested reader to [Sat09, §§5–6], in which the log geometric approach is taken for fantastacks.

Logical Dependence of Sections

The logical dependence of sections is roughly as follows.



Acknowledgments

We thank Jesse Kass and Martin Olsson for conversations which helped get this project started. We would also like to thank Vera Serganova and the MathOverflow community (especially Torsten Ekedahl, Jim Humphreys, Peter McNamara, David Speyer, and Angelo Vistoli) for their help with several technical points. Finally, we would like to thank Smiley for helping to track down many references.

2 Definitions

For a brief introduction to algebraic stacks, we refer the reader to [Ols08, Chapter 1]. For a more detailed treatment, we refer to [Vis05] or [LMB00]. If the reader is unfamiliar with stacks, we encourage her to continue reading, simply treating $[X/G]$ as a formal quotient of a scheme X by an action of a group G. Just as it is possible to learn the theory of toric varieties as a means of learning about varieties in general, we hope the theory of toric stacks can serve as an introduction to the theory of algebraic stacks.

We refer the reader to [Ful93, Chapter 1] or [CLS11, Chapter 3] for the standard correspondence between fans on lattices and toric varieties, and for basic results about toric varieties. We will follow the notation in [CLS11] whenever possible.

Definition 2.1. If L is a lattice, we denote by T_L the torus $D(L^*) = \mathrm{Hom}_{\mathrm{gp}}(\mathrm{Hom}_{\mathrm{gp}}(L,\mathbb{Z}),\mathbb{G}_m)$ whose lattice of 1-parameter subgroups is naturally isomorphic to L.

Remark 2.2. Here, the functor $D(-)$ is the *Cartier dual* $\mathrm{Hom}_{\mathrm{gp}}(-,\mathbb{G}_m)$. It is an anti-equivalence of categories between finitely generated abelian groups and diagonalizable group schemes. See [SGA3, Exposé VIII].

The functor $(-)^*$ is the usual dual for finitely generated abelian groups, $\mathrm{Hom}_{\mathrm{gp}}(-,\mathbb{Z})$.

2.1 The Toric Stack of a Stacky Fan

Suppose X is a toric variety and $G \subseteq T_0$ is a subgroup of its torus. We may then encode the toric stack $[X/G]$ combinatorially as follows.

Associated to the toric variety X is a fan Σ on the lattice of 1-parameter subgroups of T_0, $L = \mathrm{Hom}_{\mathrm{gp}}(\mathbb{G}_m, T_0)$ (see [Ful93, §1.4] or [CLS11, §3.1]). The surjection of tori $T_0 \to T_0/G$ is encoded by the induced homomorphism of lattices of 1-parameter subgroups, $\beta\colon L \to N = \mathrm{Hom}_{\mathrm{gp}}(\mathbb{G}_m, T_0/G)$. The dual homomorphism, $\beta^*\colon \mathrm{Hom}(N,\mathbb{Z}) \to \mathrm{Hom}(L,\mathbb{Z})$, is the induced homomorphism of characters. Since $T_0 \to T_0/G$ is surjective, β^* is injective, so the image of β has finite index (c.f. Remark 2.16).

Definition 2.3. A *stacky fan* is a pair (Σ,β), where Σ is a fan on a lattice L, and $\beta\colon L \to N$ is a homomorphism to a lattice N such that $\beta(L)$ has finite index in N.

The above paragraph has shown that any toric stack can be encoded with a stacky fan. Conversely, any stacky fan (Σ,β) gives rise to a toric stack as follows. Let X_Σ be the toric variety associated to Σ. The dual of β, $\beta^*\colon N^* \to L^*$, induces a homomorphism of tori $T_\beta\colon T_L \to T_N$, naturally identifying β with the induced map on lattices of 1-parameter subgroups. Since $\beta(L)$ is finite index in N, β^* is injective, so T_β is surjective. Let $G_\beta = \ker(T_\beta)$. Note that T_L is the torus of X_Σ, and $G_\beta \subseteq T_L$ is a subgroup.

Definition 2.4. Using the notation in the above paragraph, if (Σ,β) is a stacky fan, we define the toric stack $\mathcal{X}_{\Sigma,\beta}$ to be $[X_\Sigma/G_\beta]$, with the torus $T_N = T_L/G_\beta$.

Example	2.5	2.6	2.8	2.9	2.11	2.12
Σ	Σ					
$\begin{tikzcd} L \arrow[d, "\beta"] \\ N \end{tikzcd}$	$\begin{tikzcd} N \arrow[d, "\mathrm{id}"] \\ N \end{tikzcd}$	$\begin{tikzcd} \mathbb{Z}^2 \arrow[d, "\left(\begin{smallmatrix}1&0\\1&2\end{smallmatrix}\right)"] \\ \mathbb{Z}^2 \end{tikzcd}$	$\begin{tikzcd} \mathbb{Z}^2 \arrow[d, "(1\ 0)"] \\ \mathbb{Z} \end{tikzcd}$	$\begin{tikzcd} \mathbb{Z}^2 \arrow[d, "(1\ \text{-}1)"] \\ \mathbb{Z} \end{tikzcd}$	$\begin{tikzcd} \mathbb{Z}^2 \arrow[d, "(1\ 1)"] \\ \mathbb{Z} \end{tikzcd}$	$\begin{tikzcd} \mathbb{Z} \arrow[d, "2"] \\ \mathbb{Z} \end{tikzcd}$

Example 2.5 (Toric Varieties). Suppose Σ is a fan on a lattice N. Letting $L = N$ and $\beta = \mathrm{id}_N$, we see that the induced map $T_N \to T_N$ is the identity map, so G_β is trivial. So $\mathcal{X}_{\Sigma,\beta}$ is the toric variety X_Σ. ◇

Example 2.6. Here $X_\Sigma = \mathbb{A}^2$. We have that β^* is given by $\left(\begin{smallmatrix}1&1\\0&2\end{smallmatrix}\right) : \mathbb{Z}^2 \to \mathbb{Z}^2$, so the induced map on tori is $\mathbb{G}_m^2 \to \mathbb{G}_m^2$ given by $(s,t) \mapsto (st, t^2)$. The kernel is $G_\beta = \mu_2 = \{(\zeta,\zeta) | \zeta^2 = 1\} \subseteq \mathbb{G}_m^2$.



So we see that $\mathcal{X}_{\Sigma,\beta} = [\mathbb{A}^2/\mu_2]$, where the action of μ_2 is given by $\zeta \cdot (x,y) = (\zeta x, \zeta y)$. Note that this is a smooth stack. It is distinct from the singular toric variety with the fan shown to the left. ◇

Since quotients of subschemes of $\mathbb{A}^n$ by subgroups of $\mathbb{G}_m^n$ appear frequently, we often include the weights of the action in the notation.

Notation 2.7. Let $G \hookrightarrow \mathbb{G}_m^n$ be the subgroup corresponding to the surjection $\mathbb{Z}^n \to D(G)$. Let g_i be the image of e_i in $D(G)$. Let $X \subseteq \mathbb{A}^n$ be a $\mathbb{G}_m^n$-invariant subscheme. We denote the quotient $[X/G]$ by $[X/_{(\,g_1\ \cdots\ g_n\,)}G]$.

In this notation, the stack in Example 2.6 would be denoted $[\mathbb{A}^2/_{(1\ 1)}\mu_2]$.

Example 2.8. Again we have that $X_\Sigma = \mathbb{A}^2$. This time $\beta^* = \left(\begin{smallmatrix}1\\0\end{smallmatrix}\right) : \mathbb{Z} \to \mathbb{Z}^2$, which induces the homomorphism $\mathbb{G}_m^2 \to \mathbb{G}_m$ given by $(s,t) \mapsto s$. Therefore, $G_\beta = \mathbb{G}_m = \{(1,t)\} \subseteq \mathbb{G}_m^2$, so $\mathcal{X}_{\Sigma,\beta} = [\mathbb{A}^2/_{(0\ 1)}\mathbb{G}_m] \cong \mathbb{A}^1 \times [\mathbb{A}^1/\mathbb{G}_m]$. ◇

Example 2.9. This time $X_\Sigma = \mathbb{A}^2 \smallsetminus \{(0,0)\}$. We see that $\beta^* = \left(\begin{smallmatrix}1\\-1\end{smallmatrix}\right) : \mathbb{Z} \to \mathbb{Z}^2$, which induces the morphism $\mathbb{G}_m^2 \to \mathbb{G}_m$ given by $(s,t) \mapsto st^{-1}$. So $G_\beta = \mathbb{G}_m = \{(t,t)\} \subseteq \mathbb{G}_m^2$. We then have that $\mathcal{X}_{\Sigma,\beta} = [(\mathbb{A}^2 \smallsetminus \{(0,0)\})/_{(1\ 1)}\mathbb{G}_m] = \mathbb{P}^1$. ◇

Warning 2.10. Examples 2.5 and 2.9 show that non-isomorphic stacky fans (see Definition 3.2) can give rise to isomorphic toric stacks. The two presentations $[(\mathbb{A}^2 \smallsetminus \{(0,0)\})/_{(1\ 1)}\mathbb{G}_m]$ and $[\mathbb{P}^1/\{e\}]$ of the same toric stack are produced by different stacky fans. In Appendix B we collect several results which mitigate this problem.

Example 2.11 (The non-separated line). Again we have that $X_\Sigma = \mathbb{A}^2 \smallsetminus \{(0,0)\}$. However, this time we see that $\beta^* = \left(\begin{smallmatrix}1\\1\end{smallmatrix}\right) : \mathbb{Z} \to \mathbb{Z}^2$, which induces the homomorphism $\mathbb{G}_m^2 \to \mathbb{G}_m$ given by $(s,t) \mapsto st$. Therefore, $G_\beta = \mathbb{G}_m = \{(t,t^{-1})\} \subseteq \mathbb{G}_m^2$. So we have that $\mathcal{X}_{\Sigma,\beta} = [(\mathbb{A}^2 \smallsetminus \{(0,0)\})/_{(1\ \text{-}1)}\mathbb{G}_m]$ is the affine line $\mathbb{A}^1$ with a doubled origin.

This example shows that there are toric stacks which are schemes, but are not toric varieties because they are non-separated. ◇

Example 2.12. Here $X_\Sigma = \mathbb{A}^1$, and $\beta^* = 2\colon \mathbb{Z} \to \mathbb{Z}$, which induces the map $\mathbb{G}_m \to \mathbb{G}_m$ given by $t \mapsto t^2$. So $G_\beta = \mu_2 \subseteq \mathbb{G}_m$ and $\mathcal{X}_{\Sigma,\beta} = [\mathbb{A}^1/\mu_2]$. ◇

2.2 The Generically Stacky Case

Before we generalize Definitions 2.3 and 2.4 to include the generically stacky case, it is convenient to establish some terminology.

Definition 2.13. Suppose B is a finitely generated abelian group and $A \subseteq B$ is a subgroup. The *saturation of A in B* is the subgroup

$$\operatorname{Sat}_B A = \{b \in B | n \cdot b \in A \text{ for some } n \in \mathbb{Z}_{>0}\}.$$

We say A is *saturated in* B if $A = \operatorname{Sat}_B A$. We say that a homomorphism $f\colon A \to B$ is *saturated* if $f(A)$ is saturated in B.

Remark 2.14. Saturated morphisms are precisely morphisms whose cokernels are lattices (free abelian groups). In particular, the image of a saturated morphism has a direct complement.

Definition 2.15. A homomorphism of finitely generated abelian groups $f\colon A \to B$ is *close* if $\operatorname{Sat}_B f(A) = B$.

Remark 2.16. Equivalently, f is close if the dual homomorphism $f^*\colon B^* = \operatorname{Hom}(B,\mathbb{Z}) \to \operatorname{Hom}(A,\mathbb{Z}) = A^*$ is injective. Note that the property of being a close morphism depends only on the quasi-isomorphism class of the mapping cone $C(f) = [A \xrightarrow{f} B]$ in the derived category of abelian groups.

Definition 2.17. Suppose $f\colon A \to B$ is a homomorphism of finitely generated abelian groups so that $\ker f$ is free. For $i = 0, 1$, let $D(G^i_f)$ be $H^i(C(f)^*)$, where $C(f) = [A \xrightarrow{f} B]$ is the mapping cone of f and $(-)^*$ is the derived functor $R\operatorname{Hom}_{\mathrm{gp}}(-,\mathbb{Z})$. We define G^i_f to be the diagonalizable groups corresponding to $D(G^i_f)$, and we define $G_f = G^0_f \oplus G^1_f$.

Note that the homomorphism $A^* \to D(G^1_f)$ induces a homomorphism $G_f \to D(A^*)$ (that is trivial on G^0_f). In the case where A and B are free abelian groups, $H^i(C(f)^*)$ are simply the kernel ($i = 0$) and cokernel ($i = 1$) of f^*. If we additionally assume f is close (as was the case in §2.1), then f^* has no kernel, so G^0_f is trivial. In particular, the notation is consistent with the notation in the paragraph above Definition 2.4, with $f = \beta$.

We now generalize Definitions 2.3 and 2.4.

Definition 2.18. A *generically stacky fan* is a pair (Σ, β), where Σ is a fan on a lattice L, and $\beta\colon L \to N$ is a homomorphism to a finitely generated abelian group.

Definition 2.19. If (Σ, β) is a generically stacky fan, we define $\mathcal{X}_{\Sigma,\beta}$ to be $[X_\Sigma/G_\beta]$, where the action of G_β on X_Σ is induced by the homomorphism $G_\beta \to D(L^*) = T_L$.

Remark 2.20. If β is assumed to be close, this construction of $\mathcal{X}_{\Sigma,\beta}$ essentially agrees with the ones in [BCS05, §3] and [Sat09, §5]. However, those constructions effectively impose additional conditions on Σ (e.g. that Σ is a subfan of the fan of $\mathbb{A}^n$) since it is required to be induced by a fan on $N \otimes_{\mathbb{Z}} \mathbb{Q}$.

Remark 2.21. Now we give a more explicit description of $\mathcal{X}_{\Sigma,\beta}$, which also has the benefit of demonstrating that it is a generically stacky toric stack according to Definition 1.1. See Example 4.15 for an illustration of this approach.

Let $(\Sigma, \beta\colon L \to N)$ be a generically stacky fan. Let

$$\mathbb{Z}^s \xrightarrow{Q} \mathbb{Z}^r \to N \to 0$$

be a presentation of N, and let $B\colon L \to \mathbb{Z}^r$ be a lift of β.

Define the fan Σ' on $L \oplus \mathbb{Z}^s$ as follows. Let τ be the cone generated by $e_1, \dots, e_s \in \mathbb{Z}^s$. For each $\sigma \in \Sigma$, let σ' be the cone spanned by σ and τ in $L \oplus \mathbb{Z}^s$. Let Σ' be the fan generated by all the σ'. Corresponding to the cone τ, we have the closed subvariety $Y \subseteq X_{\Sigma'}$, which isomorphic to X_Σ since Σ is the *star* (sometimes called the *link*) of τ [CLS11, Proposition 3.2.7]. We define

$$\begin{aligned} \beta' = B \oplus Q\colon\ L \oplus \mathbb{Z}^s &\longrightarrow \mathbb{Z}^r \\ (l, \mathbf{a}) &\longmapsto B(l) + Q(\mathbf{a}). \end{aligned}$$

Then (Σ', β') is a stacky fan and we see that $\mathcal{X}_{\Sigma,\beta} \cong [Y/G_{\beta'}]$. Note that $C(\beta')$ is quasi-isomorphic to $C(\beta)$, so $G_{\beta'} \cong G_\beta$.

Remark 2.22. Note that if σ is a smooth cone,[2] then the cone spanned by σ and τ is also a smooth cone. So if $\mathcal{X}_{\Sigma,\beta}$ is a *smooth* generically stacky toric stack, then it is a closed substack of a *smooth* toric stack.

Remark 2.23 (On the condition "β is close"). Since the action of G^0_β on X_σ is trivial, we have that $\mathcal{X}_{\Sigma,\beta} = [X_\Sigma/G^1_\beta] \times BG^0_\beta$. It is often easiest to treat this extra stackiness separately. Let N_1 be the saturation of $\beta(L)$ in N, let N_0 be a direct complement, and let $\beta_1\colon L \to N_1$ be the factorization of β through N_1. Then $G^0_\beta = D(N_0^*) = \mathbb{G}_m^{\mathrm{rk}(N_0)}$ and $[X_\Sigma/G^1_\beta] = \mathcal{X}_{\Sigma,\beta_1}$.

We therefore typically assume β is close (or equivalently that $\mathcal{X}_{\Sigma,\beta}$ has finite generic stabilizer), with the understanding that the non-close case can usually be handled by replacing β by β_1.

Remark 2.24 (On the generically stacky case). In this paper, we opt to work primarily with toric stacks, since generically stacky toric stacks can be described as closed substacks.

The primary reason for this focus is that we would like to avoid discussing stacky tori and their actions. We refer the interested reader to [FMN09, §1.7, §2, and Appendix B] for a discussion on stacky tori and their actions. All stacky tori that arise in generically stacky toric stacks are of the form $T \times BG$, where T is a (non-stacky) torus, and G is a diagonalizable group.

However, we do deal with generically stacky fans whenever it is possible to do so without delving too heavily into the theory of stacky tori.

3 Morphisms of Toric Stacks

The main goal of this section is to define morphisms of toric stacks and stacky fans, and to show (in Theorem 3.4) that every morphism of toric stacks is induced by a morphism of stacky fans.

---

[2] A *smooth cone* is a cone whose corresponding toric variety is smooth. See [CLS11, Definition 1.2.16].

Definition 3.1. A *toric morphism* or a *morphism of (generically stacky) toric stacks* is a morphism which restricts to a homomorphism of (stacky) tori.

Definition 3.2. A *morphism of generically stacky fans* $(\Sigma, \beta : L \to N) \to (\Sigma', \beta' : L' \to N')$ is a pair of group morphisms $\Phi\colon L \to L'$ and $\phi\colon N \to N'$ so that $\beta' \circ \Phi = \phi \circ \beta$ and so that for every cone $\sigma \in \Sigma$, $\Phi(\sigma)$ is contained in a cone of Σ'.

We typically draw a morphism of generically stacky fans as a commutative diagram.

$$\begin{tikzcd}
\Sigma \arrow[r] & \Sigma' \\
L \arrow[r, "\Phi"] \arrow[d, "\beta"'] & L' \arrow[d, "\beta'"] \\
N \arrow[r, "\phi"] & N'
\end{tikzcd}$$

A morphism of generically stacky fans $(\Phi, \phi)\colon (\Sigma, \beta) \to (\Sigma', \beta')$ induces a morphism of toric varieties $X_\Sigma \to X_{\Sigma'}$ and a compatible morphism of groups $G_\beta \to G_{\beta'}$, so it induces a toric morphism of (generically stacky) toric stacks $\mathcal{X}_{(\Phi,\phi)}\colon \mathcal{X}_{\Sigma,\beta} \to \mathcal{X}_{\Sigma',\beta'}$.

3.1 Toric Morphisms are Induced by Morphisms of Stacky Fans

Lemma 3.3. Let X be a connected scheme, G a group scheme over k, and $P \to X$ a G-torsor. Suppose $Q \subseteq P$ is a connected component of P.[3] Then $Q \to X$ is an H-torsor, where H is the subgroup of G which sends Q to itself.

Proof. Let ϕ be an automorphism of P. Since Q is a connected component of P, $\phi(Q)$ is either equal to Q or is disjoint from Q. It follows that $(G \times Q) \times_P Q = (H \times Q) \times_P Q \cong H \times Q$, where the map $G \times Q \to P$ is induced by the action of G on P.

The diagonal $Q \to Q \times_X Q \subseteq P \times_X Q$ is a section of the G-torsor $P \times_X Q \to Q$, so it induces a G-equivariant isomorphism $P \times_X Q \cong G \times Q$. We then have the following cartesian diagram.

$$\begin{tikzcd}
H \times Q \cong (G \times Q) \times_P Q \arrow[r] \arrow[d] & Q \times_X Q \arrow[r] \arrow[d] & Q \arrow[d] \\
G \times Q \arrow[r, phantom, "\cong"] & P \times_X Q \arrow[r] \arrow[d] & P \arrow[d] \\
 & Q \arrow[r] & X
\end{tikzcd}$$

In particular, the map $H \times Q \to Q \times_X Q$, given by $(h, q) \mapsto (h \cdot q, q)$, is an isomorphism. This shows that Q is an H-torsor. $\square$

Theorem 3.4. *Let $(\Sigma, \beta\colon L \to N)$ and $(\Sigma', \beta'\colon L' \to N')$ be stacky fans, and suppose $f\colon \mathcal{X}_{\Sigma,\beta} \to \mathcal{X}_{\Sigma',\beta'}$ is a toric morphism. Then there exists a stacky fan (Σ_0, β_0) and morphisms $(\Phi, \phi)\colon (\Sigma_0, \beta_0) \to$*

---

[3] The hypothesis that X is connected is actually unnecessary, but it makes the condition on Q simpler. It is enough to assume that every connected component of X has exactly one connected component of Q lying over it.

(Σ, β) *and* $(\Phi', \phi')\colon (\Sigma_0, \beta_0) \to (\Sigma', \beta')$ *such that the following triangle commutes and* $\mathcal{X}_{(\Phi,\phi)}$ *is an isomorphism.*

$$\begin{tikzcd}
\mathcal{X}_{\Sigma_0,\beta_0} \arrow[dr, "\mathcal{X}_{(\Phi',\phi')}"] \arrow[dd, "\mathcal{X}_{(\Phi,\phi)}"', "\wr"] & \\
& \mathcal{X}_{\Sigma',\beta'} \\
\mathcal{X}_{\Sigma,\beta} \arrow[ur, "f"'] &
\end{tikzcd}$$

Proof. By assumption, f restricts to a homomorphism of tori $T_N \to T_{N'}$, which induces a homomorphism of lattices of 1-parameter subgroups $\phi\colon N \to N'$.

We define $Y = X_\Sigma \times_{\mathcal{X}_{\Sigma',\beta'}} X_{\Sigma'}$. Since $X_\Sigma \to \mathcal{X}_{\Sigma',\beta'}$ and $X_{\Sigma'} \to \mathcal{X}_{\Sigma',\beta'}$ are toric, $T_L \times_{T_{N'}} T_{L'}$ is a diagonalizable group. The connected component of the identity, $T_0 \subseteq T_L \times_{T_{N'}} T_{L'}$, is a connected diagonalizable group, so it is a torus. Let Y_0 be the connected component of Y which contains T_0, and let G_Φ be the kernel of the homomorphism $T_0 \to T_L$. We then have the following diagram.

$$\begin{tikzcd}
G_\Phi \arrow[rr, hook] \arrow[d, hook] & & G_{\beta'} \arrow[rr, equal] \arrow[d, hook] & & G_{\beta'} \arrow[d, hook] & \\
T_0 \arrow[rr, hook] \arrow[dr, hook] & & T_L \times_{T_{N'}} T_{L'} \arrow[rr] \arrow[dr, hook] \arrow[dd, two heads] & & T_{L'} \arrow[dr, hook] \arrow[dd, two heads] & \\
& Y_0 \arrow[rr, hook] & & Y \arrow[rr] \arrow[dd] & & X_{\Sigma'} \arrow[dd, "G_{\beta'}\text{-torsor}"] \\
& & T_L \arrow[rr] \arrow[dr, hook] & & T_{N'} \arrow[dr, hook] & \\
& & & X_\Sigma \arrow[rr] & & \mathcal{X}_{\Sigma',\beta'}
\end{tikzcd}$$

Since X_Σ is normal and separated, and Y is a $G_{\beta'}$-torsor over X_Σ, we have that Y is normal and separated, so Y_0 is normal, separated, and connected. In particular, Y_0 is irreducible. We have that T_0 is an open subscheme of Y_0, and T_0 acts on Y_0 in a way that extends the multiplication, so Y_0 is a toric variety with torus T_0. Say it corresponds to a fan Σ_0 on the lattice L_0 of 1-parameter subgroups of T_0.

Now $Y_0 \to X_\Sigma$ and $Y_0 \to X_{\Sigma'}$ are morphisms of toric varieties, so they are induced by morphisms of fans $\Sigma_0 \to \Sigma$ and $\Sigma_0 \to \Sigma'$. Defining β_0 to be the composition $L_0 \xrightarrow{\Phi} L \xrightarrow{\beta} N$, we have morphisms of stacky fans

$$\begin{tikzcd}
\Sigma & \Sigma_0 \arrow[l] \arrow[r] & \Sigma' \\
L \arrow[d, "\beta"'] & L_0 \arrow[l, "\Phi"'] \arrow[r] \arrow[d, "\beta_0"'] & L' \arrow[d, "\beta'"] \\
N \arrow[r, equal] & N \arrow[r, "\phi"] & N'
\end{tikzcd}$$

Note that G_Φ is the kernel of the surjection $T_0 \to T_L$, so the notation is consistent with Definition 2.17. By construction, G_Φ is the subgroup of $G_{\beta'}$ which takes T_0 to itself, so it is the subgroup which takes $Y_0 = \overline{T_0}$ to itself. By Lemma 3.3, Y_0 is a G_Φ-torsor over X_Σ.

Since T_0 is a connected component of a group that surjects onto T_L, the induced morphism $T_0 \to T_L$ is surjective, so Φ is close. By Lemma A.2, G_{β_0} is an extension of G_β by G_Φ. The morphism of stacky fans $(\Sigma_0, \beta_0) \to (\Sigma, \beta)$ induces the isomorphism $\mathcal{X}_{\Sigma_0,\beta_0} = [Y/G_{\beta_0}] \to [(Y/G_\Phi)/G_\beta] = \mathcal{X}_{\Sigma,\beta}$.

On the other hand, the morphism $(\Sigma_0, \beta_0) \to (\Sigma', \beta')$ induces the morphism $[Y_0/G_{\beta_0}] \cong [X_\Sigma/G_\beta] \to \mathcal{X}_{\Sigma',\beta'}$. ☐

We conclude this section with the following proposition, which is useful for understanding when a morphism of generically stacky fans induces a toric morphism with a given property.

Proposition 3.5. *Let* $\mathbf{P}$ *be a property of morphisms which is stable under composition and base change. For* $i = 0, 1$*, let* $(\Phi_i, \phi_i)\colon (\Sigma_i, \beta_i\colon L_i \to N_i) \to (\Sigma_i, \beta_i\colon L_i \to N_i)$ *is a morphism of generically stacky fans. Then the product morphism* $(\Phi_0 \times \Phi_1, \phi_0 \times \phi_1)$ *induces an morphism of generically stacky toric stacks which has property* $\mathbf{P}$ *if and only if each* (Φ_i, ϕ_i) *does.*

$$\begin{tikzcd}
\Sigma_0 \times \Sigma_1 \arrow[r] & \Sigma_0' \times \Sigma_1' \\
L_0 \times L_1 \arrow[r, "\Phi_0\times\Phi_1"] \arrow[d, "\beta_0\times\beta_1"'] & L_0' \times L_1' \arrow[d, "\beta_0'\times\beta_1'"] \\
N_0 \times N_1 \arrow[r, "\phi_0\times\phi_1"] & N_0' \times N_1'
\end{tikzcd}$$

(see [CLS11, Proposition 3.1.14] for basic facts about product fans)

Proof. Products of morphisms with property $\mathbf{P}$ have property $\mathbf{P}$ because $f \times g = (f \times \mathrm{id}) \circ (\mathrm{id} \times g)$, and $f \times \mathrm{id}$ (resp. $\mathrm{id} \times g$) is a base change of f (resp. g).

The constructions of $\mathcal{X}_{\Sigma,\beta}$ from (Σ, β) and of $\mathcal{X}_{(\Phi,\phi)}$ from (Φ, ϕ) commute with products, so $(\Phi_0 \times \Phi_1, \phi_0 \times \phi_1)$ induces the product morphism $\mathcal{X}_{(\Phi_0\times\Phi_1,\phi_0\times\phi_1)} = \mathcal{X}_{(\Phi_0,\phi_0)} \times \mathcal{X}_{(\Phi_1,\phi_1)}$. It follows that if $\mathcal{X}_{(\Phi_i,\phi_i)}$ have property $\mathbf{P}$, then so does $\mathcal{X}_{(\Phi_0\times\Phi_1,\phi_0\times\phi_1)}$.

Conversely, suppose $\mathcal{X}_{(\Phi_0\times\Phi_1,\phi_0\times\phi_1)} = \mathcal{X}_{(\Phi_0,\phi_0)} \times \mathcal{X}_{(\Phi_1,\phi_1)}$ has property $\mathbf{P}$. We have that $\mathcal{X}_{\Sigma_0',\beta_0'}$ has a k-point, induced by the identity element of its torus, which induces a morphism $\mathcal{X}_{\Sigma_1',\beta_1'} \to \mathcal{X}_{\Sigma_0',\beta_0'} \times \mathcal{X}_{\Sigma_1',\beta_1'}$. Base changing $\mathcal{X}_{(\Phi_0,\phi_0)} \times \mathcal{X}_{(\Phi_1,\phi_1)}$ by this morphism, we get $\mathcal{X}_{(\Phi_1,\phi_1)}$, so $\mathcal{X}_{(\Phi_1,\phi_1)}$ has property $\mathbf{P}$. Similarly, $\mathcal{X}_{(\Phi_0,\phi_0)}$ has property $\mathbf{P}$. ☐

4 Fantastacks: Easy-to-Draw Examples

In this section, we introduce a broad class of smooth toric stacks which are especially easy to handle because N is a lattice and the fan on L is induced by a fan on N.

Definition 4.1. Let Σ be a fan on a lattice N, and $\beta\colon \mathbb{Z}^n \to N$ a close homomorphism so that every ray of Σ contains some $\beta(e_i)$ and every $\beta(e_i)$ lies in the support of Σ. For a cone $\sigma \in \Sigma$, let $\hat{\sigma} = \mathit{cone}(\{e_i | \beta(e_i) \in \sigma\})$. We define the fan $\widehat{\Sigma}$ on $\mathbb{Z}^n$ as the fan generated by all the $\hat{\sigma}$. We define $\mathcal{F}_{\Sigma,\beta} = \mathcal{X}_{\widehat{\Sigma},\beta}$. Any toric stack isomorphic to some $\mathcal{F}_{\Sigma,\beta}$ is called a *fantastack*.

Remark 4.2. The cones of $\widehat{\Sigma}$ are indexed by sets $\{e_{i_1}, \dots, e_{i_k}\}$ such that $\{\beta(e_{i_1}), \dots, \beta(e_{i_k})\}$ is contained in a single cone of Σ. It is therefore easy to identify which open subvariety of $\mathbb{A}^n$ is represented by $\widehat{\Sigma}$. Explicitly, define the ideal

$$J_\Sigma = \Big(\prod_{\beta(e_i) \notin \sigma} x_i \;\Big|\; \sigma \in \Sigma \Big).$$

(Note, as in the Cox construction of a toric variety, that J_Σ is generated by the monomials $\prod_{\beta(e_i)\notin\sigma} x_i$, where σ varies over *maximal* cones of Σ.) Then $X_{\widehat{\Sigma}} = \mathbb{A}^n \smallsetminus V(J_\Sigma)$.

Remark 4.3. Since β is a homomorphism of lattices, $C(\beta)^*$ can be computed by simply dualizing β. Since β is assumed to be close, $G^0_\beta = 0$, so $G_\beta = D(\operatorname{cok}\beta^*)$.

If $f\colon \mathbb{Z}^n \to \operatorname{cok}\beta^*$ is the cokernel of β^*, with $g_i = f(e_i)$, then we have that $\mathcal{F}_{\Sigma,\beta} = \big[(\mathbb{A}^n \smallsetminus V(J_\Sigma))/_{(\,g_1\ \cdots\ g_n\,)}D(\operatorname{cok}\beta^*)\big]$ using Notation 2.7.

Remark 4.4. Fantastacks are precisely the toric Artin stacks in [Sat09] which have trivial generic stabilizers.

Remark 4.5. The fantastack $\mathcal{F}_{\Sigma,\beta}$ has the toric variety X_Σ as its good moduli space, as we will show in Example 6.19 (this is also proved in [Sat09, Theorem 5.5]). In fact, a smooth toric stack $\mathcal{X}$ is a fantastack if and only if it has a toric variety X as a good moduli space and the morphism $\mathcal{X} \to X$ restricts to an isomorphism of tori.

Example 4.6. Let $N = 0$ and Σ the trivial fan on N. Let $\beta\colon \mathbb{Z}^n \to N$ be the zero map. Then $\widehat{\Sigma}$ is the fan of $\mathbb{A}^n$, and $G_\beta = \mathbb{G}_m^n$, so $\mathcal{F}_{\Sigma,\beta} = [\mathbb{A}^n/\mathbb{G}_m^n]$. ◇

Example 4.7. By Remark B.22, any smooth toric variety is a fantastack. If X_Σ is a smooth toric variety, then we construct $\beta\colon \mathbb{Z}^n \to \Sigma$ by sending the generators of $\mathbb{Z}^n$ to the first lattice points along the rays of Σ. Then $X_\Sigma \cong \mathcal{F}_{\Sigma,\beta}$.

For a general (non-smooth) fan Σ, one can still construct β as above, but the resulting fantastack $\mathcal{F}_{\Sigma,\beta}$ is not isomorphic to X_Σ. However, it is the canonical stack over X_Σ, a sort of minimal stacky resolution of singularities. See Section 5. ◇

Notation 4.8. When describing fantastacks, we draw the fan Σ and label $\beta(e_i)$ with the number i.



Example 4.9



Example 4.10



Example 4.11



Example 4.12

Example 4.9. Since a single cone contains all the $\beta(e_i)$, we have that $X_{\widehat{\Sigma}} = \mathbb{A}^2$ (see Remark 4.2). We have $\beta = (1\ 1)\colon \mathbb{Z}^2 \to \mathbb{Z}$, so we compute the cokernel

$$\mathbb{Z} \xrightarrow{\beta^*=\left(\begin{smallmatrix}1\\1\end{smallmatrix}\right)} \mathbb{Z}^2 \xrightarrow{(\,1\ \text{-}1\,)} \mathbb{Z}.$$

We see that $\mathcal{F}_{\Sigma,\beta} = [\mathbb{A}^2/_{(\,1\ \text{-}1\,)}\mathbb{G}_m]$ (see Remark 4.3). ◇

Example 4.10. Since a single cone contains all the $\beta(e_i)$, we have that $X_{\widehat{\Sigma}} = \mathbb{A}^2$ (c.f. Remark 4.2). We compute the cokernel of β^*

$$\mathbb{Z}^2 \xrightarrow{\beta^*=\left(\begin{smallmatrix}1&0\\1&2\end{smallmatrix}\right)} \mathbb{Z}^2 \xrightarrow{(\,1\ 1\,)} \mathbb{Z}/2.$$

Note that β^* can be read off of the picture directly: the rows of β^* are simply the coordinates of the $\beta(e_i)$.

Therefore, $\mathcal{F}_{\Sigma,\beta} = [\mathbb{A}^2/_{(1\ 1)}\mu_2]$ (c.f. Remark 4.3). This is a "stacky resolution" of the A_1 singularity

$$\begin{aligned}\mathbb{A}^2/_{(1\ 1)}\mu_2 &= \operatorname{Spec}(k[x_1,x_2]^{\mu_2})\\ &= \operatorname{Spec} k[x_1^2, x_1x_2, x_2^2] = \operatorname{Spec}\bigl(k[x,y,z]/(xy-z^2)\bigr).\end{aligned}$$

◇

Example 4.11. As in the previous examples, $X_{\widehat{\Sigma}}$ is all of $\mathbb{A}^3$. We compute the cokernel of β^*

$$\mathbb{Z}^2 \xrightarrow{\beta^* = \begin{pmatrix} 1 & 0\\ 1 & 1\\ 0 & 1\end{pmatrix}} \mathbb{Z}^3 \xrightarrow{(1\ -1\ 1)} \mathbb{Z}.$$

So $\mathcal{F}_{\Sigma,\beta} = [\mathbb{A}^3/_{(1\ \text{-}1\ 1)}\mathbb{G}_m]$.

Note that refining the fan yields an *open* substack. In this example, consider what happens when we refine the fan Σ to the fan Σ' below.



Here G_β is unchanged; indeed, G_β depends only on β, not on Σ. However, we remove the cone $cone(e_1,e_2)$ from $\widehat{\Sigma}'$. The resulting stack is therefore the open substack $\mathcal{F}_{\Sigma',\beta} = [(\mathbb{A}^3 \smallsetminus V(x_1,x_2))/_{(1\ -1\ 1)}\mathbb{G}_m]$, which is the blowup of $\mathbb{A}^2$ at the origin (c.f. Example 4.7).

The birational transformation $Bl_0(\mathbb{A}^2) \to \mathbb{A}^2$ can therefore be realized as the morphism of good moduli spaces induced by the open immersion $\mathcal{F}_{\Sigma',\beta} \to \mathcal{F}_{\Sigma,\beta}$. ◇

Example 4.12. We have that $X_{\widehat{\Sigma}} = \mathbb{A}^4$. We compute the cokernel

$$\mathbb{Z}^3 \xrightarrow{\beta^* = \begin{pmatrix} 0 & 0 & 1\\ 1 & 0 & 1\\ 1 & 1 & 1\\ 0 & 1 & 1\end{pmatrix}} \mathbb{Z}^4 \xrightarrow{(1\ \text{-}1\ 1\ \text{-}1)} \mathbb{Z},$$

so $\mathcal{F}_{\Sigma,\beta} = [\mathbb{A}^4/_{(1\ \text{-}1\ 1\ \text{-}1)}\mathbb{G}_m]$. We will see in Section 5 that $\mathcal{F}_{\Sigma,\beta}$ is the canonical stack over the singular toric variety

$$\begin{aligned}X_\Sigma = \operatorname{Spec} k[x_1,x_2,x_3,x_4]^{\mathbb{G}_m} &= \operatorname{Spec} k[x_1x_2, x_3x_4, x_1x_4, x_2x_3]\\ &= \operatorname{Spec}\bigl(k[x,y,z,w]/(xy-zw)\bigr).\end{aligned}$$

It can be regarded as a "stacky resolution" of the singularity.

Note that the two standard toric small resolutions of this singularity are both open substacks of this stacky resolution.





The fan on the left is $[(\mathbb{A}^4 \smallsetminus V(x_2, x_4))/_{(1\ \text{-}1\ 1\ \text{-}1)}\mathbb{G}_m]$ and the fan on the right is $[(\mathbb{A}^4 \smallsetminus V(x_1, x_3))/_{(1\ \text{-}1\ 1\ \text{-}1)}\mathbb{G}_m]$. These are both toric varieties (c.f. Example 4.7). $\diamond$

4.1 Some Non-fantastack Examples

Example 4.13. Suppose $\{n_1, \ldots, n_k\}$ is a set of positive integers. Let N be $\mathbb{Z}^r \oplus \bigoplus_{i=1}^{k}(\mathbb{Z}/n_i\mathbb{Z})$, L be 0, Σ be the trivial fan on L, and $\beta\colon L \to N$ the zero map.

To compute G_β, we take a free resolution of $C(\beta)$, namely

$$\mathbb{Z}^k \xrightarrow{diag(n_1,\ldots,n_k)\oplus 0} \mathbb{Z}^k \oplus \mathbb{Z}^r.$$

Then we see that

$$H^0(C(\beta)^*) = D(G^0_\beta) = \mathbb{Z}^r$$
$$H^1(C(\beta)^*) = D(G^1_\beta) = \bigoplus_{i=1}^{k}(\mathbb{Z}/n_i\mathbb{Z})$$

Therefore, $G_\beta = \mathbb{G}_m^r \times \prod \mu_{n_i}$.

Since $X_\Sigma = \operatorname{Spec} k$, we have that $\mathcal{X}_{\Sigma,\beta} = BG_\beta$. $\diamond$

Example 4.14 (C.f. [BCS05, Examples 2.1 and 3.5]). Consider the stacky fan in which $L = \mathbb{Z}^2$, $\Sigma = \llcorner$ is the subfan of $\mathbb{A}^2$ corresponding to $\mathbb{A}^2 \smallsetminus \{0\}$, $N = \mathbb{Z} \oplus (\mathbb{Z}/2)$, and $\beta = \left(\begin{smallmatrix} 2 & \text{-}3 \\ 1 & 0 \end{smallmatrix}\right) : \mathbb{Z}^2 \to \mathbb{Z} \oplus (\mathbb{Z}/2)$. Then $C(\beta)^*$ is represented by the map $\left(\begin{smallmatrix} 2 & 1 \\ \text{-}3 & 0 \\ 0 & 2 \end{smallmatrix}\right) : \mathbb{Z}^2 \to \mathbb{Z}^3$. This map is injective, and its cokernel is $(6\ 4\ \text{-}3)\colon \mathbb{Z}^3 \to \mathbb{Z}$. Therefore, $G_\beta = \mathbb{G}_m$, and the induced map to $T_L = \mathbb{G}_m^2$ is given by $t \mapsto (t^6, t^4)$. So $\mathcal{X}_{\Sigma,\beta} = [(\mathbb{A}^2 \smallsetminus \{0\})/_{(6\ 4)}\mathbb{G}_m]$. This is the weighted projective stack $\mathbb{P}(6,4)$, or the moduli stack of elliptic curves $\overline{\mathcal{M}}_{1,1}$. $\diamond$

We repeat the previous example to illustrate that it can be realized as a closed substack of a fantastack. This approach is explained in the discussion following Definition 2.19.

Example 4.15. Consider the stacky fan in which $L = \mathbb{Z}^2$, $\Sigma = \llcorner$ is the subfan of $\mathbb{A}^2$ corresponding to $\mathbb{A}^2 \smallsetminus \{0\}$, $N = \mathbb{Z} \oplus (\mathbb{Z}/2)$, and $\beta = \left(\begin{smallmatrix} 2 & \text{-}3 \\ 1 & 0 \end{smallmatrix}\right) : \mathbb{Z}^2 \to \mathbb{Z} \oplus (\mathbb{Z}/2)$.

We replace β by the quasi-isomorphic map $\beta' = \left(\begin{smallmatrix} 2 & -3 & 0 \\ 1 & 0 & 2 \end{smallmatrix}\right) : \mathbb{Z}^3 \to \mathbb{Z}^2$, and the fan Σ by the fan Σ' obtained by adding the cone τ.


τ
$\Sigma =$
$= \Sigma'$
$L = \mathbb{Z}^2$
$L \oplus \mathbb{Z}$
β
$\beta' = \left(\begin{smallmatrix} 2 & -3 & 0 \\ 1 & 0 & 2 \end{smallmatrix}\right)$
N
$\mathbb{Z} \oplus \mathbb{Z}$



We see that $\mathcal{X}_{\Sigma',\beta'}$ is the fantastack corresponding to the fan on the right. Explicitly, it is the fantastack $[(\mathbb{A}^3 \smallsetminus V(x_1, x_2))/_{(6\ 4\ \text{-}3)}\mathbb{G}_m]$. The closed substack $\mathcal{X}_{\Sigma,\beta}$ is the divisor corresponding to the "extra ray," which is numbered 3 in the picture. That is, it is the divisor $V(x_3) = [(\mathbb{A}^2 \smallsetminus V(x_1, x_2))/_{(6\ 4)}\mathbb{G}_m]$. ◇

5 Canonical Stacks

The Cox construction demonstrates that any toric stack is the good moduli space of a smooth stack. Given a generically stacky toric stack, there is a canonical smooth generically stacky toric stack of which it is a good moduli space. The purpose of this section is to construct and characterize this canonical smooth stack.

Given a fan Σ on a lattice L, the Cox construction [CLS11, §5.1] of the toric variety X_Σ produces an open subscheme U of $\mathbb{A}^n$ and a subgroup $H \subseteq \mathbb{G}_m^n$ so that $X_\Sigma = U/H$. That is, so that $[U/H] \to X_\Sigma$ is a good moduli space in the sense of [Alp08]. We recall and generalize this construction here.

Let (Σ, β) be a generically stacky fan. Let $\Sigma(1)$ be the set of rays of Σ. Let $M \subseteq L$ be the saturated sublattice spanned by Σ, and let $M' \subseteq L$ be a direct complement to M. For each ray $\rho \in \Sigma(1)$, let u_ρ be the first element of M along ρ, and let e_ρ be the generator in $\mathbb{Z}^{\Sigma(1)}$ corresponding to ρ. We then have a morphism $\Phi\colon \mathbb{Z}^{\Sigma(1)} \times M' \to L$ given by $(e_\rho, m) \mapsto u_\rho + m$. We define a fan $\widetilde{\Sigma}$ on $\mathbb{Z}^{\Sigma(1)} \times M'$. For each $\sigma \in \Sigma$, we define $\tilde{\sigma} \in \widetilde{\Sigma}$ as the cone generated by $\{e_\rho | \rho \in \sigma\}$. The morphism of generically stacky fans

$$\begin{tikzcd}
\widetilde{\Sigma} \arrow[r] & \Sigma \\
\mathbb{Z}^{\Sigma(1)} \times M' \arrow[r, "\Phi"] \arrow[d, "\tilde{\beta}"'] & L \arrow[d, "\beta"] \\
N \arrow[r, equal] & N
\end{tikzcd}$$

induces a toric morphism $\mathcal{X}_{\widetilde{\Sigma},\tilde{\beta}} \to \mathcal{X}_{\Sigma,\beta}$.

Remark 5.1. The usual Cox construction expresses X_Σ as a quotient of $X_{\widetilde{\Sigma}}$ by G_Φ. Applying Lemma A.1 (Φ is close by construction), we see that the morphism constructed above is obtained by quotienting the morphism $[X_{\widetilde{\Sigma}}/G_\Phi] \to X_\Sigma$ by the action of G_β. This shows that the construction above commutes with quotienting $\mathcal{X}_{\Sigma,\beta}$ by its torus (i.e. replacing G_β by $G_{L\to 0}$).

Remark 5.2. Moreover, since $[\mathcal{X}_{\widetilde{\Sigma}}/G_\Phi] \to X_\Sigma$ is a good moduli space morphism, so is $\mathcal{X}_{\widetilde{\Sigma},\tilde{\beta}} \to \mathcal{X}_{\Sigma,\beta}$ by [Alp08, Proposition 4.6].

Definition 5.3. We call $\mathcal{X}_{\widetilde{\Sigma},\tilde{\beta}}$ the *canonical stack* over $\mathcal{X}_{\Sigma,\beta}$, and we say that the morphism $\mathcal{X}_{\widetilde{\Sigma},\tilde{\beta}} \to \mathcal{X}_{\Sigma,\beta}$ is a *canonical stack morphism.*

The remainder of this subsection is dedicated to justifying this terminology by showing that the canonical stack has a universal property (Proposition 5.7).

Remark 5.4. The universal property itself is basically useless. There is a more practical universal property using the language of log geometry, but a useless universal property is good enough for our needs. The only purpose of demonstrating the universal property is to show that the canonical stack is uniquely determined by the stack $\mathcal{X}_{\Sigma,\beta}$ together with the torus action (rather than by the stacky fan (Σ,β)). In particular, if a stack $\mathcal{X}$ (with a dense open torus that acts on it) has an open cover by toric stacks, the canonical stacks over the open substacks are canonically isomorphic on overlaps, so they glue to a canonical stack over $\mathcal{X}$. This will be important in the proof of [GS11b, Theorem 6.1].

Lemma 5.5. Let $\mathcal{X}_{\Sigma,\beta}$ be a cohomologically affine toric stack with n torus-invariant irreducible divisors. Suppose $f\colon \mathcal{X}_{\Sigma',\beta'} \to \mathcal{X}_{\Sigma,\beta}$ is a toric surjection from a smooth cohomologically affine toric stack with n torus-invariant divisors, which restricts to an isomorphism on tori. Then $\mathcal{X}_{\Sigma',\beta'} \to \mathcal{X}_{\Sigma,\beta}$ factors uniquely through the canonical stack over $\mathcal{X}_{\Sigma,\beta}$.

Proof. By Theorem 3.4, we may assume f is induced by a morphism of stacky fans $(\Phi,\phi)\colon (\Sigma',\beta') \to (\Sigma,\beta)$. Since f restricts to an isomorphism on tori, ϕ must be an isomorphism.

Since we are only considering torus-equivariant morphisms, we may verify the property after quotienting by the action of the torus, so we may assume $\mathcal{X}_{\Sigma,\beta}$ is the quotient of an affine toric variety by its torus, and $\mathcal{X}_{\Sigma',\beta'}$ is the quotient of a smooth affine toric variety with n-divisors by its torus. This identifies $\mathcal{X}_{\Sigma',\beta'}$ as $[\mathbb{A}^n/\mathbb{G}_m^n]$, so we may assume Σ' is the fan of $\mathbb{A}^n$. We identify the first lattice points along the rays of Σ' with the generators $e_i \in \mathbb{Z}^n$.

Since f is surjective, the induced morphism $\Sigma' \to \Sigma$ is surjective by Lemma B.3. Every ray of Σ is then the image of a unique ray of Σ', since Σ' has only n rays. Suppose $\Phi(e_i) = k_i\rho_i$. Then we see that Φ factors uniquely through the canonical stack via the morphism of fans $\Sigma' \to \widetilde{\Sigma}$ given by sending e_i to $k_i e_{\rho_i}$. ☐

Remark 5.6 ("Canonical stacks are stable under base change by open immersions"). The pre-image of a torus-invariant divisor of $\mathcal{X}_{\Sigma,\beta}$ is a divisor in its canonical stack. So by the above lemma, restricting a canonical stack morphism to the open complement of a torus-invariant divisor yields a canonical stack morphism.

As a corollary, we get the following Proposition.

Proposition 5.7 (Universal property of the canonical stack)**.** *Suppose $\mathcal{X}_{\Sigma',\beta'} \to \mathcal{X}_{\Sigma,\beta}$ is a toric morphism from a smooth toric stack, which restricts to an isomorphism of tori, and which restricts to a canonical stack morphism over every torus-invariant cohomologically affine open substack of $\mathcal{X}_{\Sigma,\beta}$. Then $\mathcal{X}_{\Sigma',\beta'}$ is canonically isomorphic to the canonical stack over $\mathcal{X}_{\Sigma,\beta}$.*

Remark 5.8. We will see in Lemma 6.7 that the morphism $\mathcal{X}_{\widetilde{\Sigma},\tilde{\beta}} \to \mathcal{X}_{\Sigma,\beta}$ is a good moduli space morphism.

By Remark B.22, we see that the canonical stack morphism over a smooth toric stack is an isomorphism. In particular, this shows that for any generically stacky toric stack $\mathcal{X}_{\Sigma,\beta}$, the canonical stack is isomorphic to $\mathcal{X}_{\Sigma,\beta}$ over its smooth locus. Thus, the canonical stack can be regarded as a (canonical!) "stacky resolution of singularities." (c.f. Examples 4.10 and 4.12)

By Remark 5.6, the definition of a canonical stack morphism can be extended to stacks which are only locally known to be toric stacks. This will be important in [GS11b], where we prove that certain stacks are toric by showing that they are locally toric, that their canonical stacks are (globally) toric, and that the property of being toric can be "descended" along canonical stack morphisms.

Definition 5.9. Suppose $\mathcal{X}$ is a stack with an open cover by generically stacky toric stacks with a common torus. A morphism from a smooth stack $\mathcal{Y} \to \mathcal{X}$ is a *canonical stack morphism* if it restricts to canonical stack morphisms on the open toric substacks of $\mathcal{X}$.

6 Toric Good Moduli Space Morphisms

6.1 Good Moduli Space Morphisms

In [Alp08], Alper introduces the notion of a good moduli space morphism, which generalizes the notion of a good quotient in geometric invariant theory and is moreover a common generalization of the notion of a tame Artin stack [AOV08] and of a coarse moduli space [KM97].

Definition 6.1. A quasi-compact and quasi-separated morphism of algebraic stacks $f\colon \mathcal{X} \to \mathcal{Y}$ is a *good moduli space morphism* if

- (f is Stein) the morphism $\mathcal{O}_{\mathcal{Y}} \to f_*\mathcal{O}_{\mathcal{X}}$ is an isomorphism, and
- (f is cohomologically affine) the pushforward functor $f_*\colon \mathsf{QCoh}(\mathcal{O}_{\mathcal{X}}) \to \mathsf{QCoh}(\mathcal{O}_{\mathcal{Y}})$ is exact.

Remark 6.2. An algebraic stack $\mathcal{X}$ is said to be cohomologically affine if the structure morphism $\mathcal{X} \to \operatorname{Spec} k$ is cohomologically affine. This is in agreement with Definition B.1. If a toric variety X_σ is an affine toric variety and G_β is an affine group, then $[X_\sigma/G_\beta] \to \operatorname{Spec} k$ is cohomologically affine by [Alp08, Proposition 3.13].

Conversely, suppose $\mathcal{X}_{\Sigma,\beta} = [X_\Sigma/G_\beta] \to \operatorname{Spec} k$ is cohomologically affine. Since X_Σ is affine over $\mathcal{X}_{\Sigma,\beta}$ (because it is a G_β-torsor), we have that $X_\Sigma \to \operatorname{Spec} k$ is cohomologically affine, so X_Σ is affine by Serre's criterion [EGA, Corollary 5.2.2]. It follows that Σ is a single cone.

Our goal in this section is to prove the following theorem which characterizes toric good moduli space morphisms:

Theorem 6.3. *Let $(\Phi,\phi)\colon (\Sigma,\beta\colon L \to N) \to (\Sigma',\beta'\colon L' \to N')$ be a morphism of generically stacky fans with β close. The induced morphism $\mathcal{X}_{\Sigma,\beta} \to \mathcal{X}_{\Sigma',\beta'}$ is a good moduli space morphism if and only if*

1. *For every $\sigma' \in \Sigma'$, the pre-image $\Phi^{-1}(\sigma')$ is a single cone $\sigma \in \Sigma$ with $\Phi(\sigma) = \sigma'$. In particular, the pre-image of the zero cone is some cone $\tau \in \Sigma$, and the pre-image of any other cone has τ as a face. Let τ^{gp} denote $(L \cap \tau)^{gp}$.*

2. τ is unstable (see Definition 6.8),

3. ϕ is surjective, and

4. the inclusion $\beta(\tau^{\mathrm{gp}}) \subseteq \ker\phi$ is close.

This result is proved in §6.2 through several lemmas. See §6.3 for examples of toric good moduli space morphisms.

6.2 Proof of Theorem 6.3

In this subsection, we gather several lemmas which help to characterize toric good moduli space morphisms. We end the subsection with a proof of Theorem 6.3.

Lemma 6.4. For $i = 0, 1$, let (Φ_i, ϕ_i) be a morphism of generically stacky fans. Then $(\Phi_0 \times \Phi_1, \phi_0 \times \phi_1)$ induces a good moduli space morphism of generically stacky toric stacks if and only if each (Φ_i, ϕ_i) does.

Proof. Good moduli spaces are stable under composition (this follows quickly from the definition) and base change [Alp08, Proposition 4.6(i)], so the result follows from Proposition 3.5. ☐

Lemma 6.5. If $f : \mathcal{X} \to \mathcal{Y}$ and $g : \mathcal{Y} \to \mathcal{Z}$ are morphisms of algebraic stacks and f is a good moduli space, then g is a good moduli space if and only if $g \circ f$ is.

Proof. As mentioned above, if g is a good moduli space, then $g \circ f$ is. For the converse, note that $g_*\mathcal{O}_{\mathcal{Y}} = g_*f_*\mathcal{O}_{\mathcal{X}} = (g \circ f)_*\mathcal{O}_{\mathcal{X}} = \mathcal{O}_{\mathcal{Z}}$, so g is Stein. To see g is cohomologically affine, first note that for any quasi-coherent sheaf $\mathcal{F}$ on $\mathcal{Y}$, we have $Rf_*f^*\mathcal{F} = \mathcal{F} \otimes Rf_*\mathcal{O}_{\mathcal{X}} = \mathcal{F}$. Therefore, $Rg_*\mathcal{F} = R(g \circ f)_*f^*\mathcal{F} = (g \circ f)_*f^*\mathcal{F} = g_*\mathcal{F}$. ☐

Throughout the rest of this subsection (until the proof of Theorem 6.3), we use the following setup. We have a morphism of generically stacky fans $(\Phi, \phi)\colon (\sigma, \beta\colon L \to N) \to (\sigma', \beta'\colon L' \to N)$, where σ is a single cone on L, and $\sigma' = \Phi(\sigma)$.

$$\begin{tikzcd} \sigma \arrow[r, two heads] & \sigma' \\ L \arrow[r, "\Phi"] \arrow[d, "\beta"'] & L' \arrow[d, "\beta'"] \\ N \arrow[r, "\phi"] & N' \end{tikzcd}$$

Lemma 6.6 ("Characterization when target is a toric variety")**.** If β is close and β' is an isomorphism, the induced map $\mathcal{X}_{\sigma,\beta} \to X_{\sigma'}$ is a good moduli space morphism if and only if every $\lambda \in N^*$ which is non-negative on σ is of the form $\phi^*(\lambda')$ for a unique $\lambda' \in N'^*$.

Proof. Since $X_\sigma = \operatorname{Spec} k[\sigma^\vee \cap L^*]$ and $X_{\sigma'} = \operatorname{Spec} k[\sigma'^\vee \cap L'^*]$ are affine and G_β is linearly reductive, the induced map $\mathcal{X}_{(\Phi,\phi)}$ is a good moduli space morphism if and only if $k[\sigma'^\vee \cap L'^*] \to k[\sigma^\vee \cap L^*]$ is the inclusion of $k[\sigma^\vee \cap L^*]^{G_\beta} = k[\sigma^\vee \cap L^*]^{G^1_\beta}$ (this last equality holds because G^0_β acts trivially on X_σ). That is, if $\pi : L^* \to D(G^1_\beta)$ is the map defining the action of G^1_β on T_L, then $\mathcal{X}_{(\Phi,\phi)}$ is a good moduli space morphism if and only if for every $\mu \in L^*$ such that $\pi(\mu) = 0$ and μ is non-negative on σ, there is a unique $\mu' \in L'^*$ which is non-negative on σ' such that $\mu = \Phi^*(\mu')$.

Suppose first that $\mathcal{X}_{(\Phi,\phi)}$ is a good moduli space morphism. If $\lambda \in N^*$ and $\beta^*(\lambda)$ is non-negative on σ, then we see there is a unique $\lambda' \in N'^*$ such that $\Phi^*\beta'^*(\lambda') = \beta^*(\lambda)$. Since $\Phi^*\beta'^*(\lambda') = \beta^*\phi^*(\lambda')$ and β^* is injective (as β is close), we see $\lambda = \phi^*(\lambda')$.

Conversely, suppose that every $\lambda \in N^*$ which is non-negative on σ is of the form $\phi^*(\lambda')$ for a unique $\lambda' \in N'^*$. Choose a free resolution

$$0 \to \mathbb{Z}^s \xrightarrow{Q} \mathbb{Z}^r \to N \to 0$$

and a lift $B : L \to \mathbb{Z}^r$ of β. Then $D(G^1_\beta)$ is the cokernel of $(BQ)^*$ and π is the composition

$$L^* \to L^* \oplus (\mathbb{Z}^s)^* \to D(G^1_\beta).$$

If $\mu \in \ker\pi$, then there exists $\lambda \in (\mathbb{Z}^r)^*$ such that $\mu = B^*(\lambda)$ and $0 = Q^*(\lambda)$. Therefore, $\lambda \in N^*$ and $\mu = \beta^*(\lambda)$. So, we see that if $\mu \in L^*$, then $\pi(\mu) = 0$ if and only if $\mu = \beta^*(\lambda)$ for some $\lambda \in N^*$. Note that λ is necessarily unique as β is close. If μ is non-negative on σ, then by hypothesis, $\lambda = \phi^*(\lambda')$ for a unique $\lambda' \in N'^*$. To conclude the proof, note that since $\Phi(\sigma) = \sigma'$, an element $\mu' \in N'^*$ is non-negative on σ' if and only if $\Phi^*(\mu')$ is non-negative on σ. ☐

Lemma 6.7 ("Isomorphism on tori implies GMS")**.** Suppose ϕ is an isomorphism and β is close. Then $\mathcal{X}_{\sigma,\beta} \to \mathcal{X}_{\sigma',\beta'}$ is a good moduli space.

Proof. First we reduce to the case where Φ is close. Let L'' be the saturation of $\Phi(L)$ in L', let β'' be the restriction of β' to L'', and let σ'' be the cone σ', regarded as a fan on L''. By assumption (i.e. the case where Φ is close), the induced morphism $\mathcal{X}_{\sigma,\beta} \to \mathcal{X}_{\sigma'',\beta''}$ is a good moduli space morphism. Note that since β is close, so is β'', so by Lemma B.14, the induced morphism $\mathcal{X}_{\sigma'',\beta''} \to \mathcal{X}_{\sigma',\beta'}$ is an isomorphism. Therefore the composition $\mathcal{X}_{\sigma,\beta} \to \mathcal{X}_{\sigma',\beta'}$ is a good moduli space morphism.

Now we consider the case when Φ is close. By Lemma A.1, G_β is an extension of $G_{\beta'}$ by G_Φ. The induced map is then $[X_\sigma/G_\beta] = \big[[X_\sigma/G_\Phi]/G_{\beta'}\big] \to [X_{\sigma'}/G_{\beta'}]$. By Lemma 6.6, $[X_\sigma/G_\Phi] \to X_{\sigma'}$ is a good moduli space. Since the property of being a good moduli space can be checked locally in the smooth topology (even in the fpqc topology, [Alp08, Proposition 4.6]), $[X_\sigma/G_\beta] \to [X_{\sigma'}/G_{\beta'}]$ is a good moduli space. ☐

Definition 6.8. We say that a cone τ of a generically stacky fan $(\Sigma, \beta\colon L \to N)$ is *unstable* if every linear functional $N \to \mathbb{Z}$ which is non-negative on the image of τ vanishes on the image of τ.

Here are two equivalent formulations of Definition 6.8: a) τ is unstable if and only if the relative interior of the image of τ in the lattice N/N_{tor} contains 0, b) τ is stable (i.e. not unstable) if $\tau \cap \ker\beta$ is contained in a proper face of τ.

Lemma 6.9 ("Quotient of a GMS is a GMS")**.** Suppose β (resp. β') factors as $L \xrightarrow{\beta_1} N_0 \xrightarrow{\beta_0} N$

(resp. $L' \xrightarrow{\beta'_1} N'_0 \xrightarrow{\beta'_0} N'$). Suppose $\phi_0 \colon N_0 \to N'_0$ makes the following diagram commute.

$$\begin{tikzcd}
\sigma \arrow[r, two heads] & \sigma' \\
L \arrow[r, "\Phi"] \arrow[d, "\beta_0"'] & L' \arrow[d, "\beta'_0"] \\
N_0 \arrow[r, "\phi_0"] \arrow[d, "\beta_1"'] & N'_0 \arrow[d, "\beta'_1"] \\
N \arrow[r, "\phi"] & N'
\end{tikzcd}$$

Suppose

1. β_0 and β'_0 are close,
2. $\ker \beta_1$ and $\ker \beta'_1$ are free,
3. ϕ_0 induces an isomorphism between $\ker \beta_1$ and $\ker \beta'_1$, and
4. ϕ induces an isomorphism between $N/\operatorname{im} \beta_1$ and $N'/\operatorname{im} \beta'_1$.

Then $\mathcal{X}_{\sigma,\beta} \to \mathcal{X}_{\sigma',\beta'}$ is a good moduli space if and only if $\mathcal{X}_{\sigma,\beta_0} \to \mathcal{X}_{\sigma',\beta'_0}$ is a good moduli space.

Proof. The first two conditions, together with Lemma A.1 imply that the rows in the following diagram are exact.

$$\begin{tikzcd}
0 \arrow[r] & G_{\beta_0} \arrow[r] \arrow[d] & G_{\beta} \arrow[r] \arrow[d] & G_{\beta_1} \arrow[r] \arrow[d, "\wr"] & 0 \\
0 \arrow[r] & G_{\beta'_0} \arrow[r] & G_{\beta'} \arrow[r] & G_{\beta'_1} \arrow[r] & 0
\end{tikzcd}$$

The last two conditions imply that the induced morphism $C(\beta_1) \to C(\beta'_1)$ is a quasi-isomorphism, which implies that the rightmost vertical map is an isomorphism between G_{β_1} and $G_{\beta'_1}$. Therefore, the morphism $\mathcal{X}_{\sigma,\beta} \cong [\mathcal{X}_{\sigma,\beta_0}/G_{\beta_1}] \to [\mathcal{X}_{\sigma',\beta'_0}/G_{\beta'_1}] \cong \mathcal{X}_{\sigma',\beta'}$ is the quotient of the morphism $\mathcal{X}_{\sigma,\beta_0} \to \mathcal{X}_{\sigma',\beta'_0}$ by $G_{\beta_1} \cong G_{\beta'_1}$. Since the property of being a good moduli space can be checked locally on the base in the smooth topology [Alp08, Proposition 4.6], $\mathcal{X}_{\sigma,\beta} \to \mathcal{X}_{\sigma',\beta'}$ is a good moduli space if and only if $\mathcal{X}_{\sigma,\beta_0} \to \mathcal{X}_{\sigma',\beta'_0}$ is. ☐

Lemma 6.10 ("Removing trivial generic stackiness is a GMS")**.** Suppose $N = N' \oplus N_0$ and that β factors through N'. Let $L' = L$. Then $[X_\sigma/G_\beta] \to [X_{\sigma'}/G_{\beta'}]$ is a good moduli space.

Proof. We have that $\beta = \beta' \oplus 0 \colon L \oplus 0 \to N' \oplus N_0$, so $G_\beta = G_{\beta'} \oplus G_0$, where G_0 acts trivially on X_σ. The map $[X_\sigma/G_0] \to X_\sigma$ is then a good moduli space, so $[X_\sigma/G_\beta] = \big[[X_\sigma/G_0]/G_{\beta'}\big] \to [X_\sigma/G_{\beta'}]$ is a good moduli space. ☐

Proposition 6.11. *Let τ be an unstable face of σ, and let τ^{gp} denote $(L\cap\tau)^{\mathrm{gp}}$. Suppose $L' = L/\tau^{\mathrm{gp}}$ and $N' = N/\beta(\tau^{\mathrm{gp}})$. Then (Φ,ϕ) induces a good moduli space morphism.*

$$\begin{tikzcd}
\sigma \arrow[r, two heads] & \sigma' = \sigma/\tau \\
L \arrow[r, "\Phi"] \arrow[d, "\beta"'] & L/\tau^{\mathrm{gp}} \arrow[d] \\
N \arrow[r, "\phi"] & N/\beta(\tau^{\mathrm{gp}})
\end{tikzcd}$$

Proof. Consider the following diagram, in which the rows are exact. We define $K = \ker(\tau^{\mathrm{gp}} \to \beta(\tau^{\mathrm{gp}}))$.

$$\begin{tikzcd}
K \arrow[r, equal] \arrow[dd, hook] & K \arrow[dd, hook] & & \\
& & \sigma \arrow[r, two heads] & \sigma/\tau \\
\tau^{\mathrm{gp}} \arrow[r, hook] \arrow[d, two heads] & L \arrow[r, two heads] \arrow[d, two heads, "\beta_0"] & L/\tau^{\mathrm{gp}} \arrow[d, equal] & \\
\beta(\tau^{\mathrm{gp}}) \arrow[r, hook] \arrow[d, equal] & L/K \arrow[r, two heads] \arrow[d, "\beta_1"] & L/\tau^{\mathrm{gp}} \arrow[d, "\beta_1'"] & \\
\beta(\tau^{\mathrm{gp}}) \arrow[r, hook] & N \arrow[r, two heads] & N/\beta(\tau^{\mathrm{gp}}) &
\end{tikzcd}$$

The top right square induces a good moduli space morphism by Lemma 6.6. Indeed, if $\lambda \in (L/K)^*$ is non-negative on σ, then it is 0 on τ, as τ is unstable. Hence, λ is induced from a unique element of $(L/\tau^{gp})^*$.

Note that $\ker\beta_1 = (\ker\beta)/K$ is free since K is saturated in $\ker\beta$ by construction. Applying Lemma 6.9, completes the proof (the conditions on kernels and cokernels follow from the 5-lemma). ☐

Lemma 6.12 ("Removing finite generic stackiness is a GMS")**.** Suppose $\ker\phi = N_0$ is finite and ϕ is surjective. Let $L' = L$. Then $[X_\sigma/G_\beta] \to [X_{\sigma'}/G_{\beta'}]$ is a good moduli space.

Proof. Consider the left-hand diagram with exact rows

$$\begin{tikzcd}
0 \arrow[r] & 0 \arrow[r] \arrow[d, "\beta_0"'] & L \arrow[r, equal] \arrow[d, "\beta"'] & L \arrow[r] \arrow[d, "\beta'"] & 0 \\
0 \arrow[r] & N_0 \arrow[r] & N \arrow[r, "\phi"] & N' \arrow[r] & 0
\end{tikzcd}
\qquad
\begin{tikzcd}
0 \arrow[r] & G_{\beta_0} \arrow[r] \arrow[d] & G_\beta \arrow[r] \arrow[d] & G_{\beta'} \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & 0 \arrow[r] & T_L \arrow[r, equal] & T_L \arrow[r] & 0
\end{tikzcd}$$

Since N_0 is finite, β_0 is close. By Lemma A.2, we get the induced right-hand diagram with exact rows. In particular, the action of G_{β_0} on X_σ is trivial, so the map $[X_\sigma/G_{\beta_0}] \to X_\sigma$ is a good moduli space. Thus, the map $[X_\sigma/G_\beta] = \big[[X_\sigma/G_{\beta_0}]/G_{\beta'}\big] \to [X_\sigma/G_{\beta'}]$ is a good moduli space map. ☐

Lemma 6.13. If $f\colon G \to H$ is a morphism of algebraic groups, then the induced morphism $BG \to BH$ is a good moduli space morphism if and only if f is surjective and $K = \ker(f)$ is linearly reductive.

Proof. We have that $\operatorname{Spec} k \to BH$ is a smooth cover, so by [Alp08, Proposition 4.6], $BG \to BH$ is a good moduli space morphism if and only $BG \times_{BH} \operatorname{Spec} k = [H/G] \to \operatorname{Spec} k$ is. Since $[H/G] \cong BK \times H/\operatorname{im}(f)$, the structure map to $\operatorname{Spec} k$ is a good moduli space morphism if and only if $H = \operatorname{im}(f)$ and K is linearly reductive. $\square$

Lemma 6.14. Suppose $L = L' = 0$ and that N and N' are finite. Then $\mathcal{X}_{(\Phi,\phi)}$ is a good moduli space morphism if and only if $\phi\colon N \to N'$ is surjective.

Proof. For $\beta\colon 0 \to N$ with N finite, we have that $\mathcal{X}_{0,\beta}$ is naturally $B(D(\operatorname{Ext}^1(N, \mathbb{Z})))$. By Lemma 6.13, we have that $\mathcal{X}_{(\Phi,\phi)}$ is a good moduli space morphism if and only if ϕ induces a surjection $D(\operatorname{Ext}^1(N, \mathbb{Z})) \to D(\operatorname{Ext}^1(N', \mathbb{Z}))$, which occurs if and only if it induces an injection $\operatorname{Ext}^1(N', \mathbb{Z}) \to \operatorname{Ext}^1(N, \mathbb{Z})$. We claim that this is equivalent to surjectivity of ϕ.

Indeed, for a short exact sequence of finite abelian groups

$$0 \to A \to B \to C \to 0$$

we get a short exact sequence

$$0 \to \operatorname{Ext}^1(C, \mathbb{Z}) \to \operatorname{Ext}^1(B, \mathbb{Z}) \to \operatorname{Ext}^1(A, \mathbb{Z}) \to 0.$$

So if ϕ is surjective, the induced map $\operatorname{Ext}^1(N', \mathbb{Z}) \to \operatorname{Ext}^1(N, \mathbb{Z})$ is injective. Conversely, if ϕ is not surjective, then it has a non-trivial cokernel C. We then have that the induced map $\operatorname{Ext}^1(N', \mathbb{Z}) \to \operatorname{Ext}^1(N, \mathbb{Z})$ factors as

$$\operatorname{Ext}^1(N', \mathbb{Z}) \to \operatorname{Ext}^1(\operatorname{im}\phi, \mathbb{Z}) \to \operatorname{Ext}^1(N, \mathbb{Z})$$

and the kernel of the first map is $\operatorname{Ext}^1(C, \mathbb{Z})$, which is non-canonically isomorphic to C, so $\operatorname{Ext}^1(N', \mathbb{Z}) \to \operatorname{Ext}^1(N, \mathbb{Z})$ is not injective. $\square$

Proposition 6.15. *Let τ be the preimage cone $\Phi^{-1}(0)$. If β is close, then (Φ, ϕ) induces a good moduli space morphism if and only if*

1. *τ is unstable,*
2. *ϕ is surjective, and*
3. *the inclusion $\beta(\tau^{\mathrm{gp}}) \subseteq \ker\phi$ is close.*

Proof. Assume first that conditions 1–3 hold. To show that $\mathcal{X}_{(\Phi,\phi)}$ is a good moduli space morphism, we factor (Φ, ϕ) as follows, and show that each square induces a good moduli space morphism.

$$\begin{tikzcd}
\sigma \arrow[r, two heads] & \sigma/\tau \arrow[r, "\sim"] & \sigma/\tau \arrow[r, two heads] & \sigma' \\
L \arrow[r] \arrow[d, "\beta"'] & L/\tau^{\mathrm{gp}} \arrow[r, equal] \arrow[d] & L/\tau^{\mathrm{gp}} \arrow[r] \arrow[d] & L' \arrow[d, "\beta'"] \\
N \arrow[r] & N/\beta(\tau^{\mathrm{gp}}) \arrow[r] & N' \arrow[r, equal] & N'
\end{tikzcd}$$

By condition 1 and Proposition 6.11, the left square induces a good moduli space morphism. By condition 3 and Lemma 6.12, the middle square induces a good moduli space morphism. Lastly, condition 2 and the fact that β is close shows that $\phi \circ \beta$ is close. Hence, β' is close, so Lemma 6.7 shows that the right square induces a good moduli space morphism.

Conversely, suppose $\mathcal{X}_{(\Phi,\phi)}$ is a good moduli space morphism. Base changing $\mathcal{X}_{(\Phi,\phi)}$ by the stacky torus of $\mathcal{X}_{\sigma',\beta'}$, we see that the following map of stacky fans also induces a good moduli space morphism

$$\begin{tikzcd}
\tau \arrow[r, two heads] & 0 \\
L \arrow[r, "\Phi"] \arrow[d, "\beta"'] & L' \arrow[d, "\beta'"] \\
N \arrow[r, "\phi"] & N'
\end{tikzcd}$$

By Lemma 6.6, the right square in the following diagram induces a good moduli space morphism, and so the composite map induces a good moduli space morphism

$$\begin{tikzcd}
\tau \arrow[r, two heads] & 0 \arrow[r] & 0 \\
L \arrow[r, "\Phi"] \arrow[d, "\beta"'] & L' \arrow[r, "\beta'"] \arrow[d, "\beta'"] & N' \arrow[d, "\beta'"] \\
N \arrow[r, "\phi"] & N' \arrow[r, equal] & N'
\end{tikzcd}$$

Another application of Lemma 6.6 (to the composite) shows that every linear functional on N which is non-negative on τ must be induced from N', and is therefore 0 on τ. This shows that τ is unstable.

Consider the following factorization of (Φ, ϕ)

$$\begin{tikzcd}
\tau \arrow[r, two heads] & 0 \arrow[r] & 0 \\
L \arrow[r] \arrow[d, "\beta"'] & L/\tau^{\mathrm{gp}} \arrow[r] \arrow[d] & L' \arrow[d, "\beta'"] \\
N \arrow[r] & N/\beta(\tau^{\mathrm{gp}}) \arrow[r] & N'
\end{tikzcd}$$

Proposition 6.11 shows that the square on the left induces a good moduli space morphism. Then by Lemma 6.5, the square on the right also induces a good moduli space morphism. Thus, we may assume σ and σ' are the zero cones. In this case, we must show that ϕ is surjective with finite kernel.

The good moduli space of $\mathcal{X}_{0,\beta}$ is Cartier dual to $(\mathrm{Sat}_N\, \beta(L))/N_{\mathrm{tor}}$ by Lemma 6.6. We have that N and N' have the same rank and that ϕ induces an isomorphism modulo torsion. Choose decompositions

$$N = N_{\mathrm{tor}} \oplus N_f \quad \text{and} \quad N' = N'_0 \oplus N'_{\mathrm{tor}} \oplus N'_f$$

where N_{tor} and N'_{tor} are torsion, N_f and N'_f are free, and N'_0 is a direct compliment to $\mathrm{Sat}_{N'}\, \beta'(L')$. This can be done so that $\phi = \phi_0 \oplus \phi_f$, where

$$\phi_0 : N_{\mathrm{tor}} \to N'_0 \oplus N'_{\mathrm{tor}} \quad \text{and} \quad \phi_f : N_f \to N'_f.$$

Let $\beta = \beta_{\mathrm{tor}} \oplus \beta_f \colon L \to N_{\mathrm{tor}} \oplus N_f$ and similarly let $\beta' = \beta'_0 \oplus \beta'_f \colon L' \to (N'_0 \oplus N'_{\mathrm{tor}}) \oplus N'_f$.

By Proposition B.17, the following squares induce isomorphisms of stacky tori:

$$\begin{tikzcd}
L \arrow[r, "\Delta"] \arrow[d, "\beta"'] & L \oplus L \arrow[r, "0\oplus\mathrm{id}"] \arrow[d, "(\beta_{\mathrm{tor}}{,}\beta_f)"] & 0 \oplus L \arrow[d, "0\oplus\beta_f"] \\
N \arrow[r, equal] & N_{\mathrm{tor}} \oplus N_f \arrow[r, equal] & N_{\mathrm{tor}} \oplus N_f
\end{tikzcd}$$

Hence, we see $\mathcal{X}_{0,\beta} \cong \mathcal{X}_{0,0\to N_{\mathrm{tor}}} \times \mathcal{X}_{\beta_f}$ and similarly for $\mathcal{X}_{0,\beta'}$. Moreover, we see that $\mathcal{X}_{(\Phi,\phi)}$ is the product of morphisms $\mathcal{X}_{0,\beta_f} \to \mathcal{X}_{0,\beta'_f}$ and $\mathcal{X}_{0,0\to N_{\mathrm{tor}}} \to \mathcal{X}_{0,0\to N'_{\mathrm{tor}}\oplus N'_0}$. By Lemma 6.4, each of these morphisms must be a good moduli space morphism. The first is a morphism of (non-stacky) tori, so it must be an isomorphism. Since the latter morphism is a good moduli space, Lemma 6.13 shows that ϕ_0 is surjective, and in particular, $N'_0 = 0$. Therefore, $\phi = \phi_0 \oplus \phi_f$ is surjective with finite kernel $\ker\phi = \ker\phi_0$. ☐

We now turn to Theorem 6.3.

Proof of Theorem 6.3. For $\sigma' \in \Sigma'$, let $X_{\sigma'}$ be the open subscheme of $X_{\Sigma'}$ corresponding to σ'. The property of being a good moduli space can be checked Zarsiki locally on the base, so it is equivalent to checking that $[X_{\Phi^{-1}(\sigma')}/G_\beta] \to [X_{\sigma'}/G_{\beta'}]$ is a good moduli space morphism for each $\sigma' \in \Sigma'$. Therefore, by Proposition 6.15, the four conditions in Theorem 6.3 imply that $\mathcal{X}_{\Sigma,\beta} \to \mathcal{X}_{\Sigma',\beta'}$ is a good moduli space morphism.

Conversely, if $[X_{\Phi^{-1}(\sigma')}/G_\beta] \to [X_{\sigma'}/G_{\beta'}]$ is a good moduli space morphism, then $[X_{\Phi^{-1}(\sigma')}/G_\beta]$ is cohomologically affine since it is cohomologically affine over a cohomologically affine stack. So $\Phi^{-1}(\sigma')$ is a single cone $\sigma \in \Sigma$. Good moduli space morphisms are surjective [Alp08, Theorem 4.14(i)], so by Lemma B.3, $\Phi(\sigma) = \sigma'$. Proposition 6.15 then shows that conditions 2–4 of Theorem 6.3 hold as well. ☐

6.3 Examples

Example 6.16. Consider the morphism $[\mathbb{A}^2/_{(1\ 1)}\mathbb{G}_m] \to \operatorname{Spec} k$ induced by the following map of stacky fans



$$\begin{tikzcd}
\mathbb{Z}^2 \arrow[r] \arrow[d, "(1\ \text{-}1)"'] & 0 \arrow[d] \\
\mathbb{Z} \arrow[r] & 0
\end{tikzcd}$$

Theorem 6.3 shows that this is a good moduli space morphism. Note that the unstable cone τ (the 2-dimensional cone) corresponds to the origin in $\mathbb{A}^2$. ◇

Example 6.17. Theorem 6.3 shows that the map

$$\begin{tikzcd}
\longrightarrow & \longrightarrow \\
\mathbb{Z} \arrow[r, "2"] \arrow[d, "2"'] & \mathbb{Z} \arrow[d, "id"] \\
\mathbb{Z} \arrow[r, "id"] & \mathbb{Z}
\end{tikzcd}$$

of stacky fans induces the good moduli space morphism $[\mathbb{A}^1/\mu_2] \to \mathbb{A}^1/\mu_2 \cong \mathbb{A}^1$. Here $\tau = 0$, so it is unstable; the other conditions of Theorem 6.3 clearly hold. ◇

Example 6.18. Theorem 6.3 shows that the Cox construction of a toric variety is a good moduli space, and more generally that canonical stack morphisms are good moduli space morphisms. For example, applying Theorem 6.3 to the map





$$\begin{tikzcd}
\mathbb{Z}^2 \arrow[r, "\left(\begin{smallmatrix}1&1\\0&2\end{smallmatrix}\right)"] \arrow[d, "\left(\begin{smallmatrix}1&1\\0&2\end{smallmatrix}\right)"'] & \mathbb{Z}^2 \arrow[d, equal] \\
\mathbb{Z}^2 \arrow[r, equal] & \mathbb{Z}^2
\end{tikzcd}$$

tells us that the toric variety on the right, the A_1 singularity, is the good moduli space of $[\mathbb{A}^2/\mu_2]$. ◇

Example 6.19. As a special case of Theorem 6.3, we recover [Sat09, Theorem 5.5], which states that the fantastack $\mathcal{F}_{\Sigma,\beta}$ has the toric variety X_Σ as its good moduli space. Given a fan Σ on a lattice N, let $\beta\colon \mathbb{Z}^n \to N$ and $\widehat{\Sigma}$ be as in Definition 4.1. Then Theorem 6.3 shows that the map induced by

$$\begin{tikzcd}
\widehat{\Sigma} \arrow[r] & \Sigma \\
\mathbb{Z}^n \arrow[r, "\beta"] \arrow[d, "\beta"'] & N \arrow[d, equal] \\
N \arrow[r, equal] & N
\end{tikzcd}$$

is a good moduli space morphism, as desired. ◇

It is often useful to think about a toric stack as "sandwiched" between its canonical stack and its good moduli space (if it has one). In this way, we often regard a toric stack as a "partial good moduli space" of its canonical stack, or as a "partial stacky resolution" of its good moduli space.

Example 6.20. Consider the stacky fan (Σ, β) shown in the center below. On the left we have the stacky fan of the corresponding canonical stack (see §5). On the right we have a toric variety. By

Theorem 6.3, the two morphism of stacky fans induce good moduli space morphisms of toric stacks.



$$\begin{tikzcd}
\mathbb{Z}^2 \arrow[r, "\Phi=\left(\begin{smallmatrix}1&1\\0&2\end{smallmatrix}\right)"] \arrow[d, "\tilde{\beta}=\left(\begin{smallmatrix}1&1\\0&4\end{smallmatrix}\right)"'] & \mathbb{Z}^2 \arrow[r, "\left(\begin{smallmatrix}1&0\\0&2\end{smallmatrix}\right)"] \arrow[d, "\beta=\left(\begin{smallmatrix}1&0\\0&2\end{smallmatrix}\right)"'] & \mathbb{Z}^2 \arrow[d, "\mathrm{id}"] \\
\mathbb{Z}^2 \arrow[r, "\mathrm{id}"] & \mathbb{Z}^2 \arrow[r, "\mathrm{id}"] & \mathbb{Z}^2
\end{tikzcd}$$

We can easily see that X_Σ is the A_1 singularity $\mathbb{A}^2/_{(\,1\;1\,)}\mu_2$, and that G_β is μ_2, but it is easiest to describe the action of μ_2 on X_Σ in terms of the canonical stack.

The canonical stack is $\mathcal{X}_{\widetilde{\Sigma},\tilde{\beta}} = [\mathbb{A}^2/_{(\,1\;\text{-}1\,)}\mu_4]$. We get the induced short exact sequence (c.f. Lemma A.1)

$$\begin{tikzcd}
0 \arrow[r] & G_\Phi \arrow[r] \arrow[d, equal] & G_{\tilde{\beta}} \arrow[r] \arrow[d, equal] & G_\beta \arrow[r] \arrow[d, equal] & 0 \\
0 \arrow[r] & \mu_2 \arrow[r, "2"] & \mu_4 \arrow[r] & \mu_2 \arrow[r] & 0
\end{tikzcd}$$

We can therefore express $\mathcal{X}_{\Sigma,\beta}$ as $\big[(\mathbb{A}^2/\mu_2)\;/\;(\mu_4/\mu_2)\big]$. We can view this either as a "partial good moduli space" of $\mathcal{X}_{\widetilde{\Sigma},\tilde{\beta}} = [\mathbb{A}^2/_{(\,1\;\text{-}1\,)}\mu_4]$, or as a "partial stacky resolution" of the singular toric variety $\mathcal{X}_{\Sigma'} = \mathbb{A}^2/_{(\,1\;\text{-}1\,)}\mu_4$. ◇

Example 6.21. Here is another interesting example of a non-smooth toric stack which is not a scheme. Consider the stacky fan (Σ, β) shown in the center below. On the left we have the stacky fan of the corresponding canonical stack (see §5). On the right we have a toric variety. By Theorem 6.3, the two morphisms of stacky fans induce good moduli space morphisms of toric stacks.



$$\begin{tikzcd}
\mathbb{Z}^2 \arrow[r, "\Phi=\left(\begin{smallmatrix}1&1\\0&2\end{smallmatrix}\right)"] \arrow[d, "\tilde{\beta}=\left(\begin{smallmatrix}2&2\\0&2\end{smallmatrix}\right)"'] & \mathbb{Z}^2 \arrow[r, "\left(\begin{smallmatrix}2&0\\0&1\end{smallmatrix}\right)"] \arrow[d, "\beta=\left(\begin{smallmatrix}2&0\\0&1\end{smallmatrix}\right)"'] & \mathbb{Z}^2 \arrow[d, "\mathrm{id}"] \\
\mathbb{Z}^2 \arrow[r, "\mathrm{id}"] & \mathbb{Z}^2 \arrow[r, "\mathrm{id}"] & \mathbb{Z}^2
\end{tikzcd}$$

Like the previous example, $\mathcal{X}_{\Sigma,\beta}$ is a quotient of the A_1 singularity $X_\Sigma = \mathbb{A}^2/_{(\,1\;1\,)}\mu_2$ by an action of μ_2. The canonical stack over it is $\mathcal{X}_{\widetilde{\Sigma},\tilde{\beta}} = \big[\mathbb{A}^2/_{\left(\begin{smallmatrix}1&0\\0&1\end{smallmatrix}\right)}(\mu_2 \times \mu_2)\big]$, and it is the "partial good moduli space" $\big[(\mathbb{A}^2/_{(\,1\;1\,)}\mu_2)/((\mu_2 \times \mu_2)/_{(\,1\;1\,)}\mu_2)\big]$. The toric stack $\mathcal{X}_{\Sigma,\beta}$ has $\mathbb{A}^2$ as its good moduli space.

In Example 4.10, we constructed a stack which "resolves the A_1 singularity by introducing stackiness." In the same informal language, this example *introduces a singularity* at a smooth point of $\mathbb{A}^2$ by introducing stackiness. ◇

Example 6.22. Let $\beta\colon \mathbb{Z}^2 \to \mathbb{Z} \oplus \mathbb{Z}/2$ be the map $\beta(x,y) = (x-y,0)$. Letting ϕ be the projection, Theorem 6.3 applied to



$$\begin{tikzcd}
\mathbb{Z}^2 \arrow[r] \arrow[d, "\beta"'] & 0 \arrow[d] \\
\mathbb{Z} \oplus \mathbb{Z}/2 \arrow[r, "\phi"] & \mathbb{Z}/2
\end{tikzcd}$$

shows that $[\mathbb{A}^2/\mathbb{G}_m \times \mu_2] \to B\mu_2$ is a good moduli space morphism. Note that this map *introduces* generic stackiness. ◇

7 Moduli Interpretation of Smooth Toric Stacks

A morphism $f\colon Y \to \mathbb{P}^n$ is equivalent to the data of a line bundle $\mathcal{L} = f^*\mathcal{O}_{\mathbb{P}^n}(1)$ and a choice of $n+1$ sections $\mathcal{O}_Y^{n+1} \to \mathcal{L}$ which generate $\mathcal{L}$. Cox generalized this moduli interpretation to smooth toric varieties [Cox95], and Perroni generalized it further to smooth toric Deligne-Mumford stacks [Per08]. The main goal for this section is to generalize it further to smooth generically stacky toric stacks.

In fact, we will see (Remark 7.10) that smooth generically stacky toric stacks are precisely the moduli stacks parametrizing tuples of generalized effective Cartier divisors [BV10, Example 2.5a] satisfying any given linear relations and any given intersection relations.

Proposition 7.1 ([SGA3, Exposé VIII, Proposition 4.1])**.** *Let G be a diagonalizable group scheme, and Y a scheme. Suppose $\mathcal{A}$ is a quasi-coherent sheaf of algebras on Y, together with an action of G (i.e. a grading $\mathcal{A} = \bigoplus_{\chi \in D(G)} \mathcal{A}_\chi$). Then $\mathcal{S}pec_Y\, \mathcal{A}$ is a G-torsor if and only if*

- *$\mathcal{A}_\chi$ is a line bundle for each $\chi \in D(G)$, and*
- *the homomorphism induced by multiplication $\mathcal{A}_\chi \otimes_{\mathcal{O}_Y} \mathcal{A}_{\chi'} \to \mathcal{A}_{\chi+\chi'}$ is an isomorphism.*

Since any G-torsor is affine over Y, it is clear that any G-torsor is of this form.

Notation 7.2. Given a collection of line bundles $\mathcal{L}_1, \dots, \mathcal{L}_n \in \operatorname{Pic}(Y)$ and $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{Z}^n$, let $\mathcal{L}^{\mathbf{a}} = \mathcal{L}_1^{\otimes a_1} \otimes \cdots \otimes \mathcal{L}_n^{\otimes a_n}$.

Let $\beta\colon \mathbb{Z}^n \to N$ be the close morphism of lattices so that $G_\beta = G \subseteq \mathbb{G}_m^n$. Then we have the presentation $N^* \xrightarrow{\beta^*} \mathbb{Z}^n \xrightarrow{\phi} D(G) \to 0$. A quasi-coherent $\mathcal{O}_Y$-algebra as in Proposition 7.1 is therefore equivalent to a collection of line bundles $\mathcal{L}_1, \dots, \mathcal{L}_n \in \operatorname{Pic}(Y)$ with isomorphisms $c_\psi\colon \mathcal{O}_Y \xrightarrow{\sim} \mathcal{L}^{\beta^*(\psi)}$ for $\psi \in N^*$, such that $c_\psi \otimes c_{\psi'} = c_{\psi+\psi'}\colon \mathcal{L}^{\beta^*(\psi)+\beta^*(\psi')} \to \mathcal{O}_Y \otimes \mathcal{O}_Y \cong \mathcal{O}_Y$.

Definition 7.3. Suppose Σ is a subfan of the fan of $\mathbb{A}^n$, and $\beta\colon \mathbb{Z}^n \to N$ is a close lattice homomorphism. A *(Σ,β)-collection* on a scheme Y consists of

- an n-tuple of line bundles $(\mathcal{L}_1, \dots, \mathcal{L}_n)$,

- global sections $s_i \in H^0(Y, \mathcal{L}_i)$ so that for each point $y \in Y$, there is a cone $\sigma \in \Sigma$ so that $s_i(y) \neq 0$ for all $e_i \notin \sigma$.
- trivializations $c_\psi \colon \mathcal{O}_Y \xrightarrow{\sim} \mathcal{L}^{\beta^*(\psi)}$ for each $\psi \in N^*$, satisfying the compatibility condition $c_\psi \otimes c_{\psi'} = c_{\psi+\psi'}$.

An *isomorphism* of (Σ, β)-collections is an n-tuple of isomorphisms of line bundles respecting the associated sections and trivializations.

Remark 7.4. Note that since N^* is a free subgroup of $\mathbb{Z}^n$, it suffices to specify c_ψ where ψ varies over a basis of N^*. If specified this way, the isomorphisms do not need to satisfy any compatibility condition. Different choices of these trivializations are related by the action of the torus (see Remark 7.8), so we often suppress the trivializations.

Remark 7.5. A line bundle with section is a *generalized effective Cartier divisor.* If the section is non-zero, the vanishing locus of the section is an effective Cartier divisor. Therefore, a (Σ, β)-collection can be defined as an n-tuple of generalized effective Cartier divisors $(D_1, \dots, D_n)$ such that $\sum a_i D_i$ is linearly equivalent to zero whenever $(a_1, \dots, a_n) \in N^*$ and such that $D_{i_1} \cap \dots, D_{i_k} = \varnothing$ whenever $e_{i_1}, \dots, e_{i_k}$ do not all lie on a single cone of Σ. Here we are suppressing the trivializations as in Remark 7.4.

Remark 7.6. Given a morphism $Y' \to Y$ and a (Σ, β)-collection on Y, we may pull back the line bundles, sections, and trivializations to produce a (Σ, β)-collection on Y'. This makes the category of (Σ, β)-collections into a fibered category over the category of schemes.

Theorem 7.7 (Moduli interpretation of smooth $\mathcal{X}_{\Sigma,\beta}$)**.** *Let Σ be a subfan of the fan for $\mathbb{A}^n$, and let $\beta \colon \mathbb{Z}^n \to N$ be close. Then $\mathcal{X}_{\Sigma,\beta}$ represents the fibered category of (Σ, β)-collections.*

Proof. A morphism $f \colon Y \to [\mathbb{A}^n / G_\beta]$ consists of a G_β-torsor $P \to Y$ and a G_β-equivariant morphism $P \to \mathbb{A}^n$. By Proposition 7.1, the data of a G_β-torsor is equivalent to a $D(G_\beta)$-graded quasi-coherent sheaf of algebras $\mathcal{A}$ such that $\mathcal{A}_\chi$ is a line bundle for each $\chi \in D(G_\beta)$. A G_β-equivariant morphism $P \to \mathbb{A}^n$ is then equivalent to a homomorphism of $\mathcal{O}_T$-algebras $\bigoplus_{\mathbf{a} \in \mathbb{N}^n} \mathcal{O}_T \to \mathcal{A}$ which respects the $D(G_\beta)$-grading (the $D(G_\beta)$-grading on the former algebra is induced by the $\mathbb{Z}^n$-grading and the homomorphism $\phi \colon \mathbb{Z}^n \to D(G_\beta)$). This is equivalent to homomorphisms of $\mathcal{O}_T$-modules $s_i \colon \mathcal{O}_T \to \mathcal{A}_{\phi(e_i)}$. Under this correspondence, the vanishing locus of s_i is the pre-image $[\mathbb{A}^{n-1}_i / G_\beta]$, where $\mathbb{A}^{n-1}_i$ is the i-th coordinate hyperplane. In particular, $(\mathcal{A}_{\phi(e_1)}, \dots, \mathcal{A}_{\phi(e_n)}, s_1, \dots, s_n)$ (along with the implicit trivializations) form a (Σ, β)-collection if and only if f factors through the open substack $\mathcal{X}_{\Sigma,\beta}$.

It is straightforward to verify that the above correspondence induces an equivalence. $\square$

Remark 7.8. Carefully following the construction in the proof shows that the action of the torus is as follows. Suppose $N^* \subseteq \mathbb{Z}^n$ is the sublattice of trivialized line bundles. Then the trivializations c_ψ have natural weights of the torus $T = \mathrm{Hom}_{\mathrm{gp}}(N^*, \mathbb{G}_m)$ associated to them. T acts on the trivializations via these weights.

Remark 7.9. This moduli interpretation is stable under base change by open immersions. Suppose a morphism $Y \to \mathcal{X}_{\Sigma,\beta}$ corresponds to the (Σ, β)-collection $(\mathcal{L}_i, s_i, c_\psi)$. Let Σ' be a subfan of Σ. Then the pullback $Y' = Y \times_{\mathcal{X}_{\Sigma,\beta}} \mathcal{X}_{\Sigma',\beta}$ is the open subscheme of Y where a subset of sections may simultaneously vanish only if Σ' contains the cone spanned by the rays corresponding to those sections.

Remark 7.10. As a sort of converse to Theorem 7.7, note that any set of intersection relations among an n-tuple of generalized effective Cartier divisors (i.e. any specification of which subsets of divisors should have empty intersection) determines a subfan Σ of the fan of $\mathbb{A}^n$. Furthermore, any[4] compatible collection of trivializations determines a subgroup $N^* = \{\mathbf{a} \in \mathbb{Z}^n | \mathcal{L}^{\mathbf{a}} \text{ is trivialized}\}$. The dual of the inclusion of N^* is a close homomorphism $\beta\colon \mathbb{Z}^n \to N$. Then $\mathcal{X}_{\Sigma,\beta}$ is the moduli stack of n-tuples of generalized effective Cartier divisors with the given intersection relations and linear relations.

7.1 Examples

The simplest examples to describe are fantastacks. See Notation 4.8 and Examples 4.9–4.12 for an explanation of the notation used below.

Remark 7.11. Any smooth toric stack contains an open substack which has a toric open immersion into a fantastack. Remark 7.9 therefore allows us to understand the moduli interpretation of non-fantastack smooth toric stacks by appropriately modifying the intersection relations.

Remark 7.12. We explicitly obtain linear relations by choosing a basis for N^*. For each basis element ψ, we get a trivialization of $\mathcal{L}^{\beta^*(\psi)}$. That is, we get trivializations of the divisors whose coefficients appear in the rows of β^*.



Example 7.13



Example 7.14



Example 7.15

We follow the less formal approach to (Σ, β)-collections explained in Remark 7.5.

Example 7.13. A morphism to the leftmost stack is a choice of three generalized effective Cartier divisors D_1, D_2, D_3 such that $D_1 \cap D_2 \cap D_3 = \varnothing$ (because no cone contains all three dots), and so that $D_1 - D_3 \sim \varnothing$ and $D_2 - D_3 \sim \varnothing$ (because $\beta^* = \left(\begin{smallmatrix}1 & 0 & -1\\ 0 & 1 & -1\end{smallmatrix}\right)$; see Remark 7.12). Here, $\varnothing$ denotes the empty divisor.

In other words, it is a choice of a line bundle and three global sections that do not all vanish at any point. This is the usual description of morphisms to $\mathbb{P}^2$. ◇

Example 7.14. A morphism to the middle stack is a choice of two generalized effective Cartier divisors D_1 and D_2 so that $D_1 + D_2 \sim \varnothing$ (because $\beta^* = (1\ 1)$; see Remark 7.12). Notice two particular morphisms from $\mathbb{A}^1$ to this stack; one given by setting $D_1 = 0$ and $D_2 = \varnothing$, and another by setting $D_1 = \varnothing$ and $D_2 = 0$. Indeed, the open substack where we impose the condition $D_1 \cap D_2 = \varnothing$ is the non-separated line. ◇

---

[4] There is *one* required relationship between the intersection relations and the trivializations. Namely, if the intersection of a *single* Cartier divisor is required to be empty (i.e. if the corresponding section of the line bundle is nowhere vanishing), then the line bundle must be trivialized. That is, if the intersection relations explicitly require the section to trivialize the line bundle, then it must be trivialized.

Example 7.15. A morphism to the rightmost stack is a choice of two generalized effective Cartier divisors D_1 and D_2 so that $D_1 + D_2 \sim \varnothing$ and $2D_2 \sim \varnothing$ (because $\beta^* = \left(\begin{smallmatrix} 1 & 1 \\ 0 & 2 \end{smallmatrix}\right)$; see Remark 7.12). Since there is a single cone that contains all the dots, there is no intersection condition on D_1 and D_2. Notice that two such divisors satisfy the relation necessary to specify a map to the stack in Example 7.14, so this stack has a morphism to the stack in that example. $\diamond$

7.2 Generically Stacky Smooth Toric Stacks

In this subsection we use the moduli interpretation to show that any smooth generically stacky toric stack is an essentially trivial gerbe over a toric stack.Suppose $Y \to \mathcal{X}_{\Sigma,\beta} = [X_\Sigma/G_\beta]$ is the morphism to a smooth toric stack corresponding to the (Σ,β)-collection $(\mathcal{L}_i, s_i, c_\psi)$. It factors through the closed substack corresponding to the j-th coordinate hyperplane of $\mathbb{A}^n$ if and only if $s_j = 0$. Theorem 7.7 (together with Remark 2.22) therefore gives us the following moduli interpretation of generically stacky smooth toric stacks.

Corollary 7.16 (of Theorem 7.7)**.** The generically stacky smooth toric substack $\mathcal{Z}$ of $\mathcal{X}_{\Sigma,\beta}$ corresponding to a coordinate subspace H of $\mathbb{A}^n$ has the following moduli interpretation. Morphisms $Y \to \mathcal{Z}$ correspond to (Σ,β)-collections on Y in which we require that $s_j = 0$ if H does not contain the j-th coordinate axis.

Definition 7.17. Suppose $\mathcal{K}$ is a line bundle on a stack $\mathcal{X}$ and b is a positive integer. The *root stack* $\sqrt[b]{\mathcal{K}/\mathcal{X}}$ is defined as the fiber product in the following diagram, where the map $\mathcal{X} \to B\mathbb{G}_m$ is the one induced by $\mathcal{K}$.

$$\begin{tikzcd}
\sqrt[b]{\mathcal{K}/\mathcal{X}} \arrow[r] \arrow[d] & B\mathbb{G}_m \arrow[d, "\hat{}b"] \\
\mathcal{X} \arrow[r] & B\mathbb{G}_m
\end{tikzcd}$$

The map $\hat{}b\colon B\mathbb{G}_m \to B\mathbb{G}_m$ is given by sending a line bundle to its b-th tensor power, induced by the group homomorphism $\mathbb{G}_m \to \mathbb{G}_m$ given by $t \mapsto t^b$.

If $\underline{\mathcal{K}} = (\mathcal{K}_1, \dots, \mathcal{K}_r)$ is an r-tuple of line bundles and $\mathbf{b} = (b_1, \dots, b_r)$ is an r-tuple of positive integers, we similarly define $\sqrt[\mathbf{b}]{\underline{\mathcal{K}}/\mathcal{X}}$ as $\mathcal{X} \times_{B\mathbb{G}_m^r} B\mathbb{G}_m^r$, where the map $\mathcal{X} \to B\mathbb{G}_m^r$ is induced by the tuple $(\mathcal{K}_1, \dots, \mathcal{K}_r)$ and the map $\hat{}\mathbf{b}\colon B\mathbb{G}_m^r \to B\mathbb{G}_m^r$ is induced by the homomorphism $\mathbb{G}_m^r \to \mathbb{G}_m^r$ given by $(t_1, \dots, t_r) \mapsto (t_1^{b_1}, \dots, t_r^{b_r})$. It is straightforward to check that $\sqrt[\mathbf{b}]{\underline{\mathcal{K}}/\mathcal{X}}$ is the fiber product of the $\sqrt[b_i]{\mathcal{K}_i/\mathcal{X}}$ over $\mathcal{X}$.

Remark 7.18. Explicitly, a morphism from a scheme (or stack) Y to the root stack $\sqrt[\mathbf{b}]{\underline{\mathcal{K}}/\mathcal{X}}$ is a morphism $f\colon Y \to \mathcal{X}$, an r-tuple of line bundles $(\mathcal{L}_1, \dots, \mathcal{L}_r)$, and isomorphisms $\mathcal{L}_i^{\otimes b_i} \cong \mathcal{K}_i$.

Definition 7.19. We say that $\mathcal{X} \to \mathcal{Y}$ is an *essentially trivial gerbe* if $\mathcal{X}$ is of the form $B(\mathbb{G}_m^s) \times \sqrt[\mathbf{b}]{\underline{\mathcal{K}}/\mathcal{Y}}$.

Proposition 7.20. *Let $\mathcal{Z}$ be a smooth generically stacky toric stack. Then $\mathcal{Z}$ is an essentially trivial gerbe over a toric stack.*

Proof. Suppose $\mathcal{Z}$ is a closed torus-invariant substack of a toric stack $\mathcal{X}_{\Sigma,\beta}$, with Σ a subfan of the fan of $\mathbb{A}^n$ (this is possible by Remark 2.22). Let $\mathcal{D}_1, \dots, \mathcal{D}_n$ be the torus-invariant divisors of $\mathcal{X}_{\Sigma,\beta}$.

Without loss of generality, $\mathcal{Z} = \mathcal{D}_1 \cap \cdots \cap \mathcal{D}_l$. By Corollary 7.16, $\mathcal{Z}$ is the stack of (Σ, β)-collections where $s_i = 0$ for $1 \leq i \leq l$.

Let Σ' be the restriction of Σ to the sublattice $\mathbb{Z}^{n-l} \subseteq \mathbb{Z}^n$ given by the last $n-l$ coordinates, let $N'^* = \mathbb{Z}^{n-l} \cap N^*$, and let $\beta' \colon \mathbb{Z}^{n-l} \to N'$ be the dual to the inclusion. For each i between 1 and l, let b_i be the smallest positive integer (if it exists) so that $(0, \ldots, 0, b_i, 0, \ldots, 0, a_{i,l+1}, \ldots, a_{i,n}) \in N^*$.Without loss of generality, we may assume these integers exist for $1 \leq i \leq r$. Define $\mathcal{K}_i = \mathcal{L}_{l+1}^{-a_{i,l+1}} \otimes \cdots \otimes \mathcal{L}_n^{-a_{i,n}}$.

Suppose $(\mathcal{L}_i, s_i, c_\psi)$ is a (Σ, β)-collection on a scheme such that $s_i = 0$ for $1 \leq i \leq l$. Then the last $n-l$ line bundles with sections form a (Σ', β')-collection, the line bundles $\mathcal{L}_i$ for $r < i \leq l$ satisfy no relations, and for $1 \leq i \leq r$, we have isomorphisms $\mathcal{L}_i^{b_i} \cong \mathcal{K}_i$. Therefore, a morphism to $\mathcal{Z}$ is precisely the data of a morphism to $B(\mathbb{G}_m^{l-r}) \times \sqrt[(b_1,\ldots,b_r)]{(\mathcal{K}_1, \ldots, \mathcal{K}_r)/\mathcal{X}_{\Sigma',\beta'}}$. □

Appendix A: Short Exact Sequences of G_βs

In this appendix, we prove two results which allow us to relate the groups of Definition 2.17. The basic advantage of expressing a group G as an extension of a quotient H by a normal subgroup N is that any quotient stack $[X/G]$ can be identified with $\big[[X/N]/H\big]$. This trick is used heavily throughout the papers.

We refer the reader to [GM96] for the relevant homological algebra.

Lemma A.1. Suppose L, L', and N' are finitely generated abelian groups. Suppose $\Phi \colon L \to L'$ is close and $\beta' \colon L' \to N'$ is a homomorphism. Suppose $\ker \Phi$ and $\ker \beta'$ are free. Then we have the following diagram, in which the rows are exact and the morphisms to $D(L^*)$ and $D(L'^*)$ are the ones described immediately after Definition 2.17.

$$\begin{tikzcd}
0 \arrow[r] & G_\Phi \arrow[r] \arrow[d, equal] & G_{\beta' \circ \Phi} \arrow[r] \arrow[d] & G_{\beta'} \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & G_\Phi \arrow[r] & D(L^*) \arrow[r] & D(L'^*) \arrow[r] & 0
\end{tikzcd}$$

Proof. By the octahedron axiom, the commutative triangle $L \to L' \to N'$ induces an exact triangle on cones $C(\Phi) \to C(\beta' \circ \Phi) \to C(\beta') \to C(\Phi)[1]$. This induces an exact triangle of duals, which induces the long exact sequence of homology groups

$$0 \to D(G^0_{\beta'}) \to D(G^0_{\beta' \circ \Phi}) \to \underbrace{D(G^0_\Phi)}_{0} \to D(G^1_{\beta'}) \to D(G^1_{\beta' \circ \Phi}) \to D(G^1_\Phi) \to 0.$$

Since Φ is close, $D(G^0_\Phi) = 0$. We therefore get the diagram with exact rows

$$\begin{tikzcd}
0 \arrow[r] & 0 \arrow[r] & G^0_{\beta' \circ \Phi} \arrow[r] & G^0_{\beta'} \arrow[r] & 0 \\
 & \oplus & \oplus & \oplus & \\
0 \arrow[r] & G^1_\Phi \arrow[r] \arrow[d, equal] & G^1_{\beta' \circ \Phi} \arrow[r] \arrow[d] & G^1_{\beta'} \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & G_\Phi \arrow[r] & D(L^*) \arrow[r] & D(L'^*) \arrow[r] & 0
\end{tikzcd}$$

□

Lemma A.2. Suppose we have the following commutative diagram, in which the rows are exact, in which L_0, L, L', N_0, N, and N' are finitely generated abelian groups, and $\ker \beta_0$, $\ker \beta$, and $\ker \beta'$ are free.

$$\begin{tikzcd}
0 \arrow[r] & L_0 \arrow[r] \arrow[d, "\beta_0"'] & L \arrow[r] \arrow[d, "\beta"'] & L' \arrow[r] \arrow[d, "\beta'"] & 0 \\
0 \arrow[r] & N_0 \arrow[r] & N \arrow[r] & N' \arrow[r] & 0
\end{tikzcd}$$

Suppose β_0 is close. Then the top row in the following diagram is exact.

$$\begin{tikzcd}
0 \arrow[r] & G_{\beta_0} \arrow[r] \arrow[d] & G_{\beta} \arrow[r] \arrow[d] & G_{\beta'} \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & D(L_0^*) \arrow[r] & D(L^*) \arrow[r] & D(L'^*) \arrow[r] & 0
\end{tikzcd}$$

Proof. We are given a short exact sequence $0 \to C(\beta_0) \to C(\beta) \to C(\beta') \to 0$, which gives us an exact triangle in the derived category. The dual is then again an exact triangle, so we get a long exact sequence of homology groups

$$0 \to D(G^0_{\beta'}) \to D(G^0_{\beta}) \to \underbrace{D(G^0_{\beta_0})}_{0} \to D(G^1_{\beta'}) \to D(G^1_{\beta}) \to D(G^1_{\beta_0}) \to 0.$$

Since β_0 is close, we have that $D(G^0_{\beta_0}) = 0$, so we get the diagram with exact rows

$$\begin{tikzcd}
0 \arrow[r] & 0 \arrow[r] & G^0_{\beta} \arrow[r] & G^0_{\beta'} \arrow[r] & 0 \\
 & \oplus & \oplus & \oplus & \\
0 \arrow[r] & G^1_{\beta_0} \arrow[r] \arrow[d] & G^1_{\beta} \arrow[r] \arrow[d] & G^1_{\beta'} \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & D(L_0^*) \arrow[r] & D(L^*) \arrow[r] & D(L'^*) \arrow[r] & 0
\end{tikzcd}$$

□

Appendix B: Non-uniqueness of Stacky Fans

B.1 The Cohomologically Affine Case

In this subsection, we study cohomologically affine generically stacky toric stacks in some detail. Roughly, the goal is to mitigate Warning 2.10 and establish a tight connection between cohomologically affine generically stacky toric stacksand their stacky fans. This is important in the proofs of some technical results, specifically Lemma 5.5 and [GS11b, Theorem 2.13].

Definition B.1. A generically stacky toric stack $\mathcal{X}_{\Sigma,\beta}$ is *cohomologically affine* if X_Σ is affine (c.f. Definition 6.1).

Recall that a toric variety is affine if and only if its fan is the set of faces of a single cone σ.

Notation B.2. As a slight abuse of notation, we use the symbol σ to denote a cone as well as the fan consisting of the cone σ and all of its faces.

Lemma B.3. Let $(\Phi, \phi)\colon (\sigma, \beta) \to (\Sigma', \beta')$ be a morphism of generically stacky fans, with σ a single cone. Suppose every torus orbit of $\mathcal{X}_{\Sigma',\beta'}$ is in the image of the induced morphism $\mathcal{X}_{(\Phi,\phi)}$ (e.g. if $\mathcal{X}_{(\Phi,\phi)}$ is surjective). Then the map $\sigma \to \Sigma'$ is a surjection. In particular, Σ' is a single cone, so $\mathcal{X}_{\Sigma',\beta'}$ is cohomologically affine.

Proof. The cones of Σ' correspond to torus orbits of $X_{\Sigma'}$, and to those of $\mathcal{X}_{\Sigma',\beta'}$. We see that every torus orbit of $X_{\Sigma'}$ is in the image of X_σ. Thus, the relative interior of every cone of Σ' contains the image of some face of σ. Therefore $\Phi(\sigma)$ intersects the relative interiors of all cones, and in particular all rays, of Σ'. Since $\Phi(\sigma)$ is a convex polyhedral cone, it follows that the $\Phi(\sigma)$ is the cone generated by the rays of Σ'. In particular, $\Sigma' \subseteq \Phi(\sigma)$. ☐

Remark B.4. The cohomological affineness condition in Lemma B.3 cannot be removed. For example, let $\Sigma =$  and $\Sigma' =$ . Then X_Σ is the blowup of $\mathbb{A}^2$ at the origin, minus the two torus-invariant points of the exceptional divisor, and $X_{\Sigma'}$ is $\mathbb{A}^2$. The natural map $X_\Sigma \to X_{\Sigma'}$ is surjective, but the map on fans is not.

Definition B.5. An affine toric variety X_σ is *pointed* if it contains a torus-invariant point. Alternatively, X_σ is pointed if σ spans the ambient lattice L. If (σ, β) is a generically stacky fan and X_σ is pointed, we say that $\mathcal{X}_{\sigma,\beta}$ is a *pointed generically stacky toric stack*. In this case, we say that (σ, β) is a *pointed generically stacky fan*.

Note that a pointed toric variety has a *unique* torus-invariant point.

Remark B.6. Note some immediate consequences of the equivalence of categories between toric varieties and fans. For any affine toric variety X_σ, there is a canonical pointed toric subvariety. Explicitly, let $L_\sigma \subseteq L$ be the sublattice spanned by σ. Then X_{σ,L_σ} is pointed, and the inclusion $L_\sigma \to L$ induces a toric closed immersion $X_{\sigma,L_\sigma} \to X_{\sigma,L}$. We have that $X_{\sigma,L}$ is (non-canonically) isomorphic to the product of X_{σ,L_σ} and the torus T_{L/L_σ}.

Remark B.7. Similarly, if $(\sigma, \beta\colon L \to N)$ is a generically stacky fan, with σ a single cone, there is a canonical morphism from a pointed toric stack. Let $L_\sigma \subseteq L$ be the saturated sublattice generated by σ, let $N_\sigma = \mathrm{Sat}_N(\beta(L_\sigma))$, and let $\beta_\sigma\colon L_\sigma \to N_\sigma$ be the morphism induced by β. Then we have a morphism of generically stacky fans $(\sigma, \beta_\sigma) \to (\sigma, \beta)$. We see that $\mathcal{X}_{\sigma,\beta}$ is (non-canonically) isomorphic to the product of the pointed generically stacky toric stack $\mathcal{X}_{\sigma,\beta_\sigma}$ and the stacky torus $\mathcal{X}_{*,L/L_\sigma \to N/N_\sigma}$, where $*$ is the trivial fan.

Remark B.8. Note that $\mathcal{X}_{\sigma,\beta_\sigma}$, as defined in Remark B.7, has the following property. Any toric morphism to $\mathcal{X}_{\sigma,\beta}$ from a pointed toric stack factors uniquely through $\mathcal{X}_{\sigma,\beta_\sigma}$. This follows immediately from the fact that any toric morphism from a pointed toric stack to a stacky torus must be trivial.

Lemma B.9. Let X_σ be a pointed affine toric variety, and let $Y \to X_\sigma$ be a toric morphism from a toric variety making Y into a G-torsor over X_σ for some group G. Then $Y \to X_\sigma$ is a trivial torsor and G is a torus. In particular, $Y \cong G \times X_\sigma$ as a toric variety, so there is a canonical toric section $X_\sigma \to Y$.

Proof. We have that G is the kernel of the homomorphism of tori induced by $Y \to X_\sigma$, so it is diagonalizable. We decompose G as a product of a finite group G_0 and a torus $\mathbb{G}_m^r$.

We then have that $Y_0 = Y/\mathbb{G}_m^r$ is a toric variety which is a G_0-torsor over X_σ. The fiber over the torus-invariant point of X_σ is then a torus invariant finite subset of Y_0. A torus has no finite-index subgroups, so any finite torus-invariant subset of a toric variety must consist of *fixed points* of the torus action. On the other hand, Y_0 is affine over X_σ, so it is affine, so it contains at most one torus fixed point. Therefore, G_0 must be trivial, so $G \cong \mathbb{G}_m^r$.

We have that G-torsors over X_σ are parametrized by $H^1(X_\sigma, G) \cong H^1(X_\sigma, \mathbb{G}_m)^r \cong \operatorname{Pic}(X_\sigma)^r$. By [CLS11, Proposition 4.2.2], we have that $\operatorname{Pic}(X_\sigma) = 0$, so all $\mathbb{G}_m^r$-torsors on X_σ are trivial. It follows that $Y = \mathbb{G}_m^r \times X_\sigma$, so by Remark B.6 there is a canonical toric section. ☐

Corollary B.10. Let $\mathcal{X}_{\sigma,\beta\colon L\to N}$ be a pointed generically stacky toric stack, where σ spans the lattice L. Let $f\colon \mathcal{X}_{\sigma,\beta} \to \mathcal{X}_{\Sigma',\beta'}$ be a homomorphism of toric stacks. Then f is induced by a morphism of stacky fans $(\sigma, \beta) \to (\Sigma', \beta')$.

Proof. Following the proof (and notation) of Theorem 3.4, we see that there is toric variety Y_0 with a toric morphism $Y_0 \to X_\sigma$ making Y_0 into a G_Φ-torsor over X_σ. By Lemma B.9, there is a canonical toric section s. This toric section induces a section of the morphism of stacky fans $(\Phi, \mathrm{id}_N)\colon (\Sigma_0, \beta_0) \to (\sigma, \beta)$.

$$\begin{tikzcd}
\sigma \arrow[r, bend left, "s"] & \Sigma_0 \arrow[l] \arrow[r] & \Sigma' \\
L \arrow[r, bend right] \arrow[d, "\beta"'] & L_0 \arrow[l, "\Phi"'] \arrow[r] \arrow[d, "\beta_0"] & L' \arrow[d, "\beta'"] \\
N \arrow[r, equal] & N \arrow[r, "\phi"] & N'
\end{tikzcd}$$

The composition $(\sigma, \beta) \to (\Sigma', \beta')$ then induces the morphism f. ☐

Corollary B.11. Let $(\Sigma, \beta\colon L \to N)$ be a generically stacky fan. There is a natural bijection between toric morphisms $\mathbb{A}^1 \to \mathcal{X}_{\Sigma,\beta}$ and elements of $\Sigma \cap L$.

Proof. This follows immediately from Corollary B.10 and the usual description of the fan of $\mathbb{A}^1$, namely $(\longrightarrow, \mathrm{id}\colon \mathbb{Z} \to \mathbb{Z})$. ☐

Lemma B.12. Let $(\Phi, \phi)\colon (\sigma, \beta\colon L \to N) \to (\sigma', \beta'\colon L' \to N')$ be a morphism of generically stacky fans, with σ and σ' single cones. Suppose the induced morphism $\mathcal{X}_{(\Phi,\phi)}$ is an isomorphism. Then Φ induces an isomorphism of monoids $\sigma \cap L \to \sigma' \cap L'$.

Proof. By Corollary B.11, elements of the monoid $\sigma \cap L$ are in bijection with toric morphisms from $\mathbb{A}^1$ to $\mathcal{X}_{\sigma,\beta}$. The isomorphism $\mathcal{X}_{(\Phi,\phi)}$ induces a bijection of these sets. On the other hand, the induced morphism is a morphism of monoids. ☐

B.2 Isomorphisms From Morphisms of (Generically) Stacky Fans

As we saw in Warning 2.10, non-isomorphic stacky fans can induce isomorphic toric stacks. In this section, we prove some useful results for identifying morphisms of stacky fans which induce isomorphisms of toric stacks.

Lemma B.13. For $i = 0, 1$, let (Φ_i, ϕ_i) be a morphism of generically stacky fans. Then $(\Phi_0 \times \Phi_1, \phi_0 \times \phi_1)$ induces an isomorphism of generically stacky toric stacks if and only if each (Φ_i, ϕ_i) does.

Proof. This is an immediate corollary of Proposition 3.5. ☐

Lemma B.14. Let Σ be a fan on a lattice L and let $\beta\colon L \to N$ be a close morphism to a finitely generated abelian group. Let L_0 be a lattice, and $\beta_0\colon L_0 \to N$ any homomorphism. Let $\Sigma\times 0$ be the fan Σ regarded as a fan on $L \oplus L_0$, supported entirely on L. Then the morphism of generically stacky fans $(\Sigma, \beta\colon L \to N) \to (\Sigma \times 0, \beta \oplus \beta_0\colon L \oplus L_0 \to N)$ induces an isomorphism $\mathcal{X}_{\Sigma,\beta} \to \mathcal{X}_{\Sigma\times 0,\beta\oplus\beta'}$.

Proof. Since $\Sigma\times 0$ is the product of Σ on L and the trivial fan on L_0, we have that $X_{\Sigma\times 0} = X_\Sigma \times T_{L_0}$.

The first diagram has exact rows.

$$\begin{tikzcd}
0 \arrow[r] & L \arrow[r] \arrow[d, "\beta"'] & L \oplus L_0 \arrow[r] \arrow[d, "\beta\oplus\beta_0"] & L_0 \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & N \arrow[r, equal] & N \arrow[r] & 0 \arrow[r] & 0
\end{tikzcd}
\qquad
\begin{tikzcd}
0 \arrow[r] & G_\beta \arrow[r] \arrow[d] & G_{\beta\oplus\beta'} \arrow[r] \arrow[d] & G_{L_0\to 0} \arrow[r] \arrow[d, "\wr"] & 0 \\
0 \arrow[r] & T_L \arrow[r] & T_L \oplus T_{L_0} \arrow[r] & T_{L_0} \arrow[r] & 0
\end{tikzcd}$$

Since β is close, Lemma A.2 implies that the second diagram has exact rows. We see that the induced morphism is then

$$\begin{aligned}
\mathcal{X}_{\Sigma,\beta} = [X_\Sigma/G_\beta] \xrightarrow{\sim} \big[[X_{\Sigma\times 0}/T_{L_0}]/G_\beta\big] &= \big[[X_{\Sigma\times 0}/G_{L_0\to 0}]/G_\beta\big] \\
&= [X_{\Sigma\times 0}/G_{\beta\oplus\beta_0}] = \mathcal{X}_{\Sigma\times 0,\beta\oplus\beta_0}.
\end{aligned}$$ ☐

Remark B.15. The condition that β is close is necessary in the above argument. As a simple counterexample where β is not close, consider the morphism of generically stacky fans $(0, \mathrm{id})\colon (\Sigma, 0 \to \mathbb{Z}) \to (\Sigma, \mathrm{id}\colon \mathbb{Z} \to \mathbb{Z})$, where Σ is the trivial fan. This induces the morphism $B\mathbb{G}_m \to \mathbb{G}_m$.

Lemma B.16. Let σ be a cone on L. Suppose $(\Phi, \phi)\colon (\sigma, \beta\colon L \to N) \to (\Phi(\sigma), \beta'\colon L' \to N')$ is a morphism of generically stacky fans so that ϕ is an isomorphism, Φ is close, and so that Φ induces an isomorphism of monoids $(\sigma \cap L) \to (\Phi(\sigma) \cap L')$. Then the induced morphism $\mathcal{X}_{\sigma,\beta} \to \mathcal{X}_{\Phi(\sigma),\beta'}$ is an isomorphism.

Proof. Let $L_\sigma \subseteq L$ be the sublattice generated by $\sigma \cap L$, and let L_1 be a direct complement. Let X_σ denote the toric variety corresponding to the cone σ, regarded as a fan on L_σ. The fan σ (on L) is the product $\sigma \times 0$ on $L_\sigma \times L_1$, so $\mathcal{X}_{\sigma,\beta} = [(X_\sigma \times T_{L_1})/G_\beta]$.

By assumption, the sublattice of L' generated by $\Phi(\sigma) \cap L'$ is isomorphic to L_σ, and the isomorphism identifies σ with $\Phi(\sigma)$. Let L'_1 be a direct complement to L_σ in L'. As above, we have that $\mathcal{X}_{\Phi(\sigma),\beta'} = [(X_\sigma \times T_{L'_1})/G_{\beta'}]$.

Note that $T_L = T_{L_\sigma} \times T_{L_1}$ and $T_{L'} = T_{L_\sigma} \times T_{L'_1}$. Since Φ is close, Lemma A.1 tells us that the following diagram has exact rows.

$$\begin{tikzcd}
0 \arrow[r] & G_\Phi \arrow[r] \arrow[d, equal] & G_\beta \arrow[r] \arrow[d] & G_{\beta'} \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & G_\Phi \arrow[r] \arrow[d, phantom, "\times"] & T_{L_1} \arrow[r] \arrow[d, phantom, "\times"] & T_{L'_1} \arrow[r] \arrow[d, phantom, "\times"] & 0 \\
0 \arrow[r] & 0 \arrow[r] & T_{L_\sigma} \arrow[r, equal] & T_{L_\sigma} \arrow[r] & 0
\end{tikzcd}$$

So we see that the toric morphism induced by (Φ, ϕ) is $\mathcal{X}_{\sigma,\beta} = [(X_\sigma \times T_{L_1})/G_\beta] = \big[[(X_\sigma \times T_{L_1})/G_\Phi]/G_{\beta'}\big] = [(X_\sigma \times T_{L_1'})/G_{\beta'}] = \mathcal{X}_{\Phi(\sigma),\beta'}$. ☐

Lemmas B.14 and B.16 can be combined and extended.

Proposition B.17. *Let $(\Phi, \phi)\colon (\sigma, \beta\colon L \to N) \to (\sigma', \beta'\colon L' \to N')$ be a morphism of generically stacky fans, with σ and σ' single cones. Suppose*

1. *ϕ is an isomorphism,*
2. *Φ induces an isomorphism of the monoids $(\sigma \cap L)$ and $(\sigma' \cap L')$, and*
3. *$\phi\big(\operatorname{Sat}_N \beta(L)\big) = \operatorname{Sat}_{N'} \beta'(L')$.*

Then the induced morphism $\mathcal{X}_{\sigma,\beta} \to \mathcal{X}_{\sigma',\beta'}$ is an isomorphism.

Proof. First we reduce to the case when β is close. Let N_0 be a direct complement to $\operatorname{Sat}_N \beta(L)$. Then we see that $(\Sigma, \beta\colon L \to N)$ is the product of $\big(\Sigma, \beta_1\colon L \to \operatorname{Sat}_N \beta(L)\big)$ and $(*, \beta_0\colon 0 \to N_0)$, where $*$ is the trivial fan (which contains only the zero cone). Condition 3 implies that $\phi(N_0)$ is a direct complement to $\operatorname{Sat}_{N'} \beta'(L')$, so the same argument shows that $(\Sigma', \beta'\colon L' \to N')$ is the product of $\big(\Sigma', \beta_1'\colon L' \to \operatorname{Sat}_{N'} \beta'(L')\big)$ and $\big(*, \beta_0'\colon 0 \to \phi(N_0)\big)$. We see that (Φ, ϕ) is a product of $\big(\Phi\colon L \to L', \phi_1\colon \operatorname{Sat}_N(\beta(L)) \to \operatorname{Sat}_{N'}(\beta'(L'))\big)$ and $(0, \phi|_{N_0})$. The later is an isomorphism by conditions 1 and 3, so by Lemma B.13, we have reduced to the case where $N = \operatorname{Sat}_N \beta(L)$.

Let L_σ be the sublattice of L generated by $\sigma \cap L$. Let N_1 be the free part of $\beta(L)$, and choose a splitting $s\colon N_1 \to L$. Let $L_1 = L_\sigma + s(N_1)$ in L.

Applying Lemmas B.16 and B.14 in succession, we see that the morphism induced by the composition $(\sigma, \beta|_{L_1}) \to (\sigma, \beta|_{\operatorname{Sat}_L L_1}) \to (\sigma, \beta)$ is an isomorphism.

Note that $\Phi|_{L_1}$ has no kernel, so we may identify $\Phi(L_1)$ with L_1. Then the same argument shows that $(\sigma, \beta|_{L_1}) \to (\sigma', \beta')$ induces an isomorphism.

We then have a factorization of morphisms of generically stacky fans $(\sigma, \beta|_{L_1}) \to (\sigma, \beta) \xrightarrow{(\Phi,\phi)} (\sigma', \beta')$. Since the first morphism induces an isomorphism and the composite induces an isomorphism, it follows that (Φ, ϕ) induces an isomorphism. ☐

Definition B.18. Suppose $\Phi\colon \Sigma \to \Sigma'$ is a morphism of fans and $\sigma' \in \Sigma'$. The *pre-image* of σ', $\Phi^{-1}(\sigma')$, is the subfan of Σ consisting of cones whose image lie in σ'.

Remark B.19. Suppose $f\colon X_\Sigma \to X_{\Sigma'}$ is the morphism of toric varieties corresponding to the map of fans $\Sigma \to \Sigma'$. The cone $\sigma' \in \Sigma'$ corresponds to an affine open subscheme $U_{\sigma'} = \operatorname{Spec}(k[\sigma'^\vee \cap N']) \subseteq X_{\Sigma'}$, the complement of the divisors corresponding to rays not on σ'. The key property of $\Phi^{-1}(\sigma')$ is that $X_\Sigma \times_{X_{\Sigma'}} U_{\sigma'}$ is naturally the open subvariety $X_{\Phi^{-1}(\sigma')} \subseteq X_\Sigma$.

Proposition B.20. *Let $(\Phi, \phi)\colon (\Sigma, \beta\colon L \to N) \to (\Sigma', \beta'\colon L' \to N')$ be a morphism of generically stacky fans. Suppose*

1. *β is close,*
2. *ϕ is an isomorphism,*

3. for every cone $\sigma' \in \Sigma'$, $\Phi^{-1}(\sigma')$ is a single cone, and

4. for every cone $\sigma' \in \Sigma'$, Φ induces a isomorphism of monoids $\Phi^{-1}(\sigma') \cap L \to \sigma' \cap L'$.

Then the induced morphism $\mathcal{X}_{\Sigma,\beta} \to \mathcal{X}_{\Sigma',\beta'}$ is an isomorphism.

Proof. We may check whether the map is an isomorphism locally on the base, so we may assume Σ' is a single cone σ'. By Proposition B.17, we get the result. □

Remark B.21. The converse of Proposition B.20 is easy to show for morphisms of toric stacks. More precisely, suppose $(\Phi, \phi)\colon (\Sigma, \beta) \to (\Sigma', \beta')$ induces an isomorphism of toric stacks. Then β is automatically close since (Σ, β) is a stacky fan (not a generically stacky fan). We have that $\mathcal{X}_{(\Phi,\phi)}$ induces an isomorphism between tori $T_N \to T_{N'}$, so the corresponding morphism of 1-parameter subgroups, $\phi\colon N \to N'$, must be an isomorphism. Since isomorphisms are affine, Remark B.19 shows that $\Phi^{-1}(\sigma')$ is a single cone for each $\sigma' \in \Sigma'$. Finally, Lemma B.12 shows that Φ induces isomorphisms of monoids.

Remark B.22. Suppose X_Σ is a smooth toric variety and $\Phi\colon (\widetilde{\Sigma}, \mathbb{Z}^n) \to (\Sigma, L)$ is its Cox construction [CLS11, §5.1]. Then the induced morphism $\mathcal{X}_{\widetilde{\Sigma},\Phi} \to \mathcal{X}_{\Sigma, id_L} = X_\Sigma$ is an isomorphism. In particular, any smooth toric variety can be expressed as a quotient of a $\mathbb{G}_m^n$-invariant open subvariety of $\mathbb{A}^n$ by a subgroup of $\mathbb{G}_m^n$. Toric stacks of this form are called *fantastacks* (see §4 and Example 4.7).

References

[Alp08] Jarod Alper. *Good moduli spaces for Artin stacks.* May 2008. http://arxiv.org/abs/0804.2242.

[AOV08] Dan Abramovich, Martin Olsson, and Angelo Vistoli. Tame stacks in positive characteristic. *Ann. Inst. Fourier (Grenoble)*, 58(4):1057–1091, 2008.

[BCS05] Lev A. Borisov, Linda Chen, and Gregory G. Smith. The orbifold Chow ring of toric Deligne-Mumford stacks. *J. Amer. Math. Soc.*, 18(1):193–215 (electronic), 2005.

[BV10] Niels Borne and Angelo Vistoli. Parabolic sheaves on logarigthmic schemes. 2010. arXiv:1001.0466.

[CLS11] David A. Cox, John B. Little, and Henry K. Schenck. *Toric Varieties*, volume 124 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2011.

[Cox95] David A. Cox. The functor of a smooth toric variety. *Tohoku Math. J. (2)*, 47(2):251–262, 1995.

[EGA] Alexander Grothendieck and Jean Dieudonné. *Éléments de géométrie algébrique.* Inst. Hautes tudes Sci. Publ. Math., 1960–1967. Nos. 4, 8, 11, 17, 20, 24, 28, 32.

[FMN09] Barbara Fantechi, Etienne Mann, and Fabio Nironi. Smooth toric DM stacks. 2009. arXiv:0708.1254v2.

[Ful93] William Fulton. *Introduction to toric varieties*, volume 131 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1993. The William H. Roever Lectures in Geometry.

[GIT] D. Mumford, J. Fogarty, and F. Kirwan. *Geometric invariant theory*, volume 34 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)]*. Springer-Verlag, Berlin, third edition, 1994.

[GM96] Sergei I. Gelfand and Yuri I. Manin. *Methods of homological algebra*. Springer-Verlag, Berlin, 1996. Translated from the 1988 Russian original.

[GS11a] Anton Geraschenko and Matthew Satriano. Toric stacks I: The theory of stacky fans. 2011. arXiv:1107.1906.

[GS11b] Anton Geraschenko and Matthew Satriano. Toric stacks II: Intrinsic characterization of toric stacks. 2011. arXiv:1107.1907.

[Iwa09] Isamu Iwanari. The category of toric stacks. *Compos. Math.*, 145(3):718–746, 2009.

[KM97] Seán Keel and Shigefumi Mori. Quotients by groupoids. *Ann. of Math. (2)*, 145(1):193–213, 1997.

[Laf02] Laurent Lafforgue. Chtoucas de Drinfeld et correspondance de Langlands. *Invent. Math.*, 147(1):1–241, 2002.

[LMB00] Gérard Laumon and Laurent Moret-Bailly. *Champs algébriques*, volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2000.

[Ols08] Martin C. Olsson. *Compactifying moduli spaces for abelian varieties*, volume 1958 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2008.

[Per08] Fabio Perroni. A note on toric Deligne-Mumford stacks. *Tohoku Math. J. (2)*, 60(3):441–458, 2008.

[Sat09] Matthew Satriano. Canonical artin stacks over log smooth schemes. 2009. arXiv:0911.2059.

[SGA3] Michel Demazure, Alexander Grothendieck, Michael Artin, J. E. Bertin, P. Gabriel, Michèle Raynaud, and J.P. Serre. *Séminaire de Géométrie Algébrique du Bois-Marie - Schémas en Groupes (SGA 3)*. Springer-Verlag, Berlin, 1970. Lecture Notes in Mathematics, Vols. 151, 152, 153.

[Tyo10] Ilya Tyomkin. Tropical geometry and correspondence theorems via toric stacks. 2010. arXiv:1001.1554v3.

[Vis05] Angelo Vistoli. *Grothendieck topologies, fibered categories, and descent theory. Part 1 of Fundamental algebraic geometry: Grothendieck's FGA explained*, volume 123 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2005.